

FORM PTO-1390 (Modified) (REV 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 29729/37581	
TRANSMITTAL LETTER TO THE UNITED STATES • DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR) 09/889075	
INTERNATIONAL APPLICATION NO. PCT/AU00/00011		INTERNATIONAL FILING DATE 11 January 2000		PRIORITY DATE CLAIMED 11 January 1999	
TITLE OF INVENTION CATALYTIC MOLECULES					
APPLICANT(S) FOR DO/EO/US DAVID G. ATKINS, ANDREW BAKER, LEVON M. KHACHIGIAN					
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:</p> <ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below. 4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c) (2)) <ol style="list-style-type: none"> a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). <ol style="list-style-type: none"> a. <input type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) <ol style="list-style-type: none"> a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)). 11. <input checked="" type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409). 12. <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210). <p>Items 13 to 20 below concern document(s) or information included:</p> <ol style="list-style-type: none"> 13. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 14. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 15. <input checked="" type="checkbox"/> A FIRST preliminary amendment. 16. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 17. <input type="checkbox"/> A substitute specification. 18. <input type="checkbox"/> A change of power of attorney and/or address letter. 19. <input checked="" type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 20. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 21. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 22. <input checked="" type="checkbox"/> Certificate of Mailing by Express Mail 23. <input type="checkbox"/> Other items or information 					

JC18 Rec'd PCT/PTO 11 JUL 2001

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.53) <div style="font-size: 1.5em; font-weight: bold;">09/889075</div>	INTERNATIONAL APPLICATION NO. <div style="font-weight: bold;">PCT/AU00/00011</div>	ATTORNEY'S DOCKET NUMBER <div style="font-weight: bold;">29729/37581</div>
-----------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------	-------------------------------------------------------------------------------

24. The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) : <input checked="" type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1000.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 <div style="text-align: right; font-weight: bold;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>	CALCULATIONS PTO USE ONLY <div style="border: 1px solid black; height: 100px; width: 100%;"></div>																																																																	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30	<div style="border: 1px solid black; padding: 2px;">\$1,000.00</div> <div style="border: 1px solid black; padding: 2px;">\$0.00</div>																																																																	
<table border="1" style="width:100%; border-collapse: collapse;"> <thead> <tr> <th style="width:15%;">CLAIMS</th> <th style="width:25%;">NUMBER FILED</th> <th style="width:25%;">NUMBER EXTRA</th> <th style="width:15%;">RATE</th> <th style="width:20%;"></th> </tr> </thead> <tbody> <tr> <td>Total claims</td> <td style="text-align: center;">43 - 20 =</td> <td style="text-align: center;">23</td> <td style="text-align: center;">x \$18.00</td> <td style="text-align: right;">\$414.00</td> </tr> <tr> <td>Independent claims</td> <td style="text-align: center;">1 - 3 =</td> <td style="text-align: center;">0</td> <td style="text-align: center;">x \$80.00</td> <td style="text-align: right;">\$0.00</td> </tr> <tr> <td colspan="4">Multiple Dependent Claims (check if applicable). <input checked="" type="checkbox"/></td> <td style="text-align: right;">\$270.00</td> </tr> <tr> <td colspan="4" style="text-align: right; font-weight: bold;">TOTAL OF ABOVE CALCULATIONS =</td> <td style="text-align: right;">\$1,684.00</td> </tr> <tr> <td colspan="4"> <input type="checkbox"/> Applicant claims small entity status. (See 37 CFR 1.27). The fees indicated above are reduced by 1/2. </td> <td style="text-align: right;">\$0.00</td> </tr> <tr> <td colspan="4" style="text-align: right; font-weight: bold;">SUBTOTAL =</td> <td style="text-align: right;">\$1,684.00</td> </tr> <tr> <td colspan="4"> Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30 + </td> <td style="text-align: right;">\$0.00</td> </tr> <tr> <td colspan="4" style="text-align: right; font-weight: bold;">TOTAL NATIONAL FEE =</td> <td style="text-align: right;">\$1,684.00</td> </tr> <tr> <td colspan="4"> Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/> </td> <td style="text-align: right;">\$0.00</td> </tr> <tr> <td colspan="4" style="text-align: right; font-weight: bold;">TOTAL FEES ENCLOSED =</td> <td style="text-align: right;">\$1,684.00</td> </tr> <tr> <td colspan="4"></td> <td style="text-align: right;">Amount to be: refunded \$</td> </tr> <tr> <td colspan="4"></td> <td style="text-align: right;">charged \$</td> </tr> </tbody></table>	CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		Total claims	43 - 20 =	23	x \$18.00	\$414.00	Independent claims	1 - 3 =	0	x \$80.00	\$0.00	Multiple Dependent Claims (check if applicable). <input checked="" type="checkbox"/>				\$270.00	TOTAL OF ABOVE CALCULATIONS =				\$1,684.00	<input type="checkbox"/> Applicant claims small entity status. (See 37 CFR 1.27). The fees indicated above are reduced by 1/2.				\$0.00	SUBTOTAL =				\$1,684.00	Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30 +				\$0.00	TOTAL NATIONAL FEE =				\$1,684.00	Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				\$0.00	TOTAL FEES ENCLOSED =				\$1,684.00					Amount to be: refunded \$					charged \$	a. <input checked="" type="checkbox"/> A check in the amount of <u>\$1,684.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>13-2855</u> A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE																																																															
Total claims	43 - 20 =	23	x \$18.00	\$414.00																																																														
Independent claims	1 - 3 =	0	x \$80.00	\$0.00																																																														
Multiple Dependent Claims (check if applicable). <input checked="" type="checkbox"/>				\$270.00																																																														
TOTAL OF ABOVE CALCULATIONS =				\$1,684.00																																																														
<input type="checkbox"/> Applicant claims small entity status. (See 37 CFR 1.27). The fees indicated above are reduced by 1/2.				\$0.00																																																														
SUBTOTAL =				\$1,684.00																																																														
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30 +				\$0.00																																																														
TOTAL NATIONAL FEE =				\$1,684.00																																																														
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				\$0.00																																																														
TOTAL FEES ENCLOSED =				\$1,684.00																																																														
				Amount to be: refunded \$																																																														
				charged \$																																																														

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

WILLIAMS, Joseph A. Jr.
 MARSHALL, O'TOOLE, GERSTEIN
 MURRAY & BORUN
 6300 Sears Tower
 233 South Wacker Drive
 Chicago, Illinois 60606-6402
 (312) 474-6300

SIGNATURE
 Joseph A. Williams, Jr.
 NAME
 38,659
 REGISTRATION NUMBER
 11 July 2001
 DATE

PCT/US 00/00011 - 2002

PATENT
30397/37581

**IN THE UNITED STATES PATENT
AND TRADEMARK OFFICE**

Applicant: **Atkins et al.**

Serial No.: 09/889,075


U.S. National Phase of PCT/AU00/00011

Filed: **January 11, 2000**

For: Catalytic Molecules

Group Art Unit: Not Yet Assigned

Examiner: Not Yet Assigned

) I hereby certify that this paper is
) being deposited with the United
) States Postal Service with sufficient
) postage as first class mail in an
) envelope addressed to:
) Commissioner for Patents,
) Washington, D.C. 20231 on this date:
)
) September 4, 2002
)
) 
) David A. Gass
) Registration No. 38,153
) Attorney for Applicants

PRELIMINARY AMENDMENT

Commissioner for Patents
 Washington, DC 20231

Sir:

Please amend the above-identified application as follows, prior to calculation of the filing fee (with extra claims) and prior to examination of the merits:

IN THE CLAIMS:

AMEND Claims 48, 49 and 59 as follows:

48. (AMENDED) A pharmaceutical composition comprising a DNAzyme according to claim 20 and a pharmaceutically acceptable carrier.

49. (AMENDED) A method of inhibiting EGR-1 activity in cells which comprises exposing the cell to a DNAzyme according to claim 20.

09/889,075

30397/37581

59. (AMENDED) An angioplastic stent for inhibition of the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactially effective dose of DNAzyme according to claim 20.

REMARKS

The applicant respectfully submits that all claims are now of proper form and scope for allowance. Early and favorable consideration is respectfully requested.

The foregoing amendments are intended solely to reduce the PTO fees for multiple dependent claims and claims in excess of 20. The applicants reserve the right to pursue subject matter of any cancelled claim in this application or any related (*e.g.*, continuing) application.

Attached hereto is a marked-up version of the changes made to the claims by the foregoing amendments. The attached page is captioned "**Version with markings to show changes made.**"

Respectfully submitted,

MARSHALL, GERSTEIN & BORUN

By: 

David A. Gass

Reg. No. 38,153

6300 Sears Tower

233 South Wacker Drive

Chicago, Illinois 60606-6402

(312) 474-6300

September 3, 2002

09/889,075

30397/37581

VERSION WITH MARKINGS TO SHOW CHANGES MADE**IN THE CLAIMS:**

AMEND Claims 48, 49 and 59 as follows:


48. (AMENDED) A pharmaceutical composition comprising a DNAzyme according to [any one of claims 20 to 47] claim 20 and a pharmaceutically acceptable carrier.
49. (AMENDED) A method of inhibiting EGR-1 activity in cells which comprises exposing the cell to a DNAzyme according to [any one of claims 20 to 47] claim 20.
59. (AMENDED) An angioplastic stent for inhibition of the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactially effective dose of DNAzyme according to [any one of claims 20 to 47] claim 20.

09/889075

JC18 Rec'd PCT/PTO 1 1 JUL 2001

PATENT
29729/37581

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: David G. Atkins et al)	“EXPRESS MAIL” mailing label No.
)	EL564462123US
U.S. National Phase of)	
International Application No:)	I hereby certify that this paper is being
PCT/AU/00/00011)	deposited with the United States Postal
filed January 11, 2000)	Service as first class mail, postage
)	prepaid, in an envelope addressed to:
Serial No: Not yet assigned)	Box PCT, Commissioner for Patents,
)	Washington, D.C. 20231, on this date
Filed: herewith)	
)	July 11, 2001
Title: CATALYTIC MOLECULES)	
)	
Group Art Unit: Not yet assigned)	
)	Richard Zimmermann
Examiner: Not yet assigned)	

PRELIMINARY AMENDMENT

Box PCT
Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

This amendment is being filed with the U.S. National filing of the above identified PCT application. The Applicant requests entry of this amendments prior to calculation of the filing fee and prior to examination on the merits.

AMENDMENTS

In the Abstract:

Please add to the application the abstract attached hereto.

In the claims:

Please cancel claims 1-19 and insert the following claims 20-62 as shown below:

20. A DNAzyme which specifically cleaves EGR-1 mRNA, the DNAzyme comprising (i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site; (ii) a first binding domain continuous with the 5' end of the catalytic domain; and (iii) a second binding domain continuous with the 3' end of the catalytic domain, wherein the binding domains are sufficiently complementary to the two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168-332 as shown in SEQ ID No: 1, such that the DNAzyme cleaves the EGR-1 mRNA.
21. A DNAzyme as claimed in claim 20 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
22. A DNAzyme as claimed in claim 20 in which the cleavage site is selected from the group consisting of
 - (i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302;
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
23. A DNAzyme as claimed in claim 22 in which the cleavage site is the AU site corresponding to nucleotides 271-272.
24. A DNAzyme as claimed in claim 22 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
25. A DNAzyme as claimed in claim 23 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
26. A DNAzyme as claimed in claim 20 in which the catalytic domain has the nucleotide sequence GGCTAGCTACAACGA [SEQ. ID. NO: 2].
27. A DNAzyme as claimed in claim 26 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.

39. A DNAzyme as claimed in claim 38 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
40. A DNAzyme as claimed in claim 38 in which the cleavage site is selected from the group consisting of
 - (i) the GU site corresponding to nucleotides 198-199;
 - (ii) the GU site corresponding to nucleotides 200-201;
 - (iii) the GU site corresponding to nucleotides 264-265;
 - (iv) the AU site corresponding to nucleotides 271-272;
 - (v) the AU site corresponding to nucleotides 301-302;
 - (vi) the GU site corresponding to nucleotides 303-304; and
 - (vii) the AU site corresponding to nucleotides 316-317.
41. A DNAzyme as claimed in claim 40 in which the cleavage site is the AU site corresponding to nucleotides 271-272.
42. A DNAzyme as claimed in claim 40 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
43. A DNAzyme as claimed in claim 41 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
44. A DNAzyme as claimed in claim 20 which has a sequence selected from the group consisting of:
 - (i) 5'-caggggacaGGCTAGCTACAACGAcgttgcg (SEQ ID NO: 3);
 - (ii) 5'-tgcaggggaGGCTAGCTACAACGAaccgttgcg (SEQ ID NO: 4);
 - (iii) 5'-catcctggaGGCTAGCTACAACGAgagcaggct (SEQ ID NO: 5);
 - (iv) 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6);
 - (v) 5'-ccgctgccaGGCTAGCTACAACGAcccgacgt (SEQ ID NO: 7);
 - (vi) 5'-gcggggacaGGCTAGCTACAACGAcagctgcat (SEQ ID NO: 8);
 - (vii) 5'-cagcggggaGGCTAGCTACAACGAatcagctgc (SEQ ID NO: 9); and
 - (viii) 5'-ggtcagagaGGCTAGCTACAACGActgcagcgg (SEQ ID NO: 10).
45. A DNAzyme as claimed in claim 44 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
46. A DNAzyme as claimed in claim 44 which has the sequence:
5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6).
47. A DNAzyme as claimed in claim 46 wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3'-end of the catalytic domain.
48. A pharmaceutical composition comprising a DNAzyme according to any one of claims 20 to 47 and a pharmaceutically acceptable carrier.

49. A method of inhibiting EGR-1 activity in cells which comprises exposing the cell to a DNAzyme according to any one of claims 20 to 47.
50. A method as claimed in claim 49 wherein the cells are vascular cells.
51. A method as claimed in any one of claims 49 wherein the cells are cells involved in neoplasia.
52. A method of inhibiting proliferation or migration of cells in a subject which comprises administering to the subject a prophylactically effective dose of the pharmaceutical composition according to claim 48.
53. A method as claimed in claim 52 wherein the cells are vascular cells.
54. A method as claimed in any one of claims 52 wherein the cells are cells involved in neoplasia.
55. A method of treating a condition associated with cell proliferation or migration in a subject which comprises administering to the subject a therapeutically effective dose of the pharmaceutical composition according to claim 48.
56. A method as claimed in claim 55 wherein the cells are vascular cells.
57. A method as claimed in any one of claims 55 wherein the cells are cells involved in neoplasia.
58. A method as claimed in claim 55 wherein the condition associated with cell proliferation or migration is selected from the group consisting of post-angioplasty restenosis, vein graft failure, hypertension, transplant coronary disease, and complications associated with atherosclerosis or peripheral vascular disease.
59. An angioplastic stent for inhibition of the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to any one of claims 20 to 47.
60. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a prophylactically effective dose of a pharmaceutical composition according to claim 48 to the subject at around the time of the angioplasty.
61. A method according to claim 60 in which the pharmaceutical composition is administered by catheter.
62. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to claim 58 to the subject at around the time of the angioplasty.

REMARKS

The Applicant is canceling the original claim set without prejudice to the Applicant's right to pursue claims of the same or similar scope in a duly filed continuing application, and is filing a new set of claims intended solely to place the claims in proper multiple dependent format. The following table demonstrates the new claims as they correspond to the original claims set:

<u>New Claim No.</u>	<u>Original Claim No. and (Dependency)</u>
Claim 20	Claim 1
Claim 21	Claim 8 (1)
Claim 22	Claim 4 (1)
Claim 23	Claim 5 (4)
Claim 24	Claim 8 (4)
Claim 25	Claim 8 (5)
Claim 26	Claim 3 (1)
Claim 27	Claim 8 (3)
Claim 28	Claim 4 (3)
Claim 29	Claim 5 (4)
Claim 30	Claim 8 (4)
Claim 31	Claim 8 (5)
Claim 32	Claim 2 (1)
Claim 33	Claim 8 (2)
Claim 34	Claim 4 (2)
Claim 35	Claim 5 (4)
Claim 36	Claim 8 (4)
Claim 37	Claim 8 (5)
Claim 38	Claim 3 (2)
Claim 39	Claim 8 (3)
Claim 40	Claim 4 (3)
Claim 41	Claim 5 (4)
Claim 42	Claim 8 (4)
Claim 43	Claim 8 (5)
Claim 44	Claim 6 (1)
Claim 45	Claim 8 (6)
Claim 46	Claim 7 (6)
Claim 47	Claim 8 (7)
Claim 48	Claim 9 (1-8)

<u>New Claim No.</u>	<u>Original Claim No. and (Dependency)</u>
Claim 49	Claim 10 (1-8)
Claim 50	Claim 13 (10)
Claim 51	Claim 14 (10)
Claim 52	Claim 11 (9)
Claim 53	Claim 13 (11)
Claim 54	Claim 14 (11)
Claim 55	Claim 12 (9)
Claim 56	Claim 13 (12)
Claim 57	Claim 14 (12)
Claim 58	Claim 15 (12)
Claim 59	Claim 16 (1-8)
Claim 60	Claim 17 (9)
Claim 61	Claim 18 (17)
Claim 62	Claim 19 (15)

The new claim set does not introduce new matter.


The abstract of the disclosure is identical to the abstract in the published PCT pamphlet.

Respectfully submitted,

MARSHALL, O'TOOLE, GERSTEIN,
MURRAY & BORUN

6300 Sears Tower
233 South Wacker Drive
Chicago, Illinois 60606

By:


Joseph A. Williams, Jr. (Reg. No. 38,659)
Attorney for Applicants

Dated: July 11, 2001

8/ARIS

09/889075
JC18 Rec'd PCT/PTO 11 JUL 2001

WO 00/42173

PCT/AU00/00011

1

CATALYTIC MOLECULES

FIELD OF THE INVENTION

The present invention relates to DNazymes which are targeted against mRNA molecules encoding EGR-1 (also known as Egr-1 or NGFI-A). The present invention also relates to compositions including these DNazymes
5 and to methods of treatment involving administration of the DNazymes.

BACKGROUND OF THE INVENTION

Egr-1 expression in Smooth Muscle Cells

Smooth muscle cells (SMCs) are well recognized as a significant
10 cellular component of atherosclerotic and post-angioplasty restenotic lesions (Stary et al, 1995; Holmes et al, 1984). SMC migration and proliferation are key events in the pathogenesis of these vascular disorders (Jackson & Schwartz, 1992; Libby et al, 1995). The promoter regions of many genes that encode mitogenic and migratory factors expressed by SMCs in these lesions
15 (Evanko et al, 1998; Murry et al, 1996; Ueda et al, 1996; Tanizawa et al, 1996; Rekhter & Gordon, 1994; Hughes et al, 1993; Brogi et al, 1993; Wilcox et al 1989; Wilcox et al, 1988) contain nucleotide (nt) recognition elements for the nuclear protein and transcription factor, Egr-1 (Khachigian & Collins, 1997; Khachigian et al, 1996). Egr-1 is not expressed in the unmanipulated artery
20 wall, but is rapidly activated by mechanical injury (Khachigian et al, 1996; Silverman et al, 1997; Kim et al, 1995). It is also induced in vascular endothelial cells and/or SMCs exposed to fluid biomechanical forces (Khachigian et al, 1997; Sumpio et al, 1998) and multiple other
25 pathophysiologically-relevant agonists (Delbridge & Khachigian, 1997).

DNazymes

In human gene therapy, antisense nucleic acid technology has been one of the major tools of choice to inactivate genes whose expression causes disease and is thus undesirable. The anti-sense approach employs a nucleic
30 acid molecule that is complementary to, and thereby hybridizes with, an mRNA molecule encoding an undesirable gene. Such hybridization leads to the inhibition of gene expression.

Anti-sense technology suffers from certain drawbacks. Anti-sense hybridization results in the formation of a DNA/target mRNA heteroduplex. This heteroduplex serves as a substrate for RNase H-mediated degradation of the target mRNA component. Here, the DNA anti-sense molecule serves in a passive manner, in that it merely facilitates the required cleavage by endogenous RNase H enzyme. This dependence on RNase H confers limitations on the design of anti-sense molecules regarding their chemistry and ability to form stable heteroduplexes with their target mRNA's. Anti-sense DNA molecules also suffer from problems associated with non-specific activity and, at higher concentrations, even toxicity.

As an alternative to anti-sense molecules, catalytic nucleic acid molecules have shown promise as therapeutic agents for suppressing gene expression, and are widely discussed in the literature (Haseloff (1988); Breaker (1994); Koizumi (1989); Otsuka; Kashani-Sabet (1992); Raillard (1996); and Carmi (1996)). Thus, unlike a conventional anti-sense molecule, a catalytic nucleic acid molecule functions by actually cleaving its target mRNA molecule instead of merely binding to it. Catalytic nucleic acid molecules can only cleave a target nucleic acid sequence if that target sequence meets certain minimum requirements. The target sequence must be complementary to the hybridizing regions of the catalytic nucleic acid, and the target must contain a specific sequence at the site of cleavage.

Catalytic RNA molecules ("ribozymes") are well documented (Haseloff (1988); Symonds (1992); and Sun (1997)), and have been shown to be capable of cleaving both RNA (Haseloff (1988)) and DNA (Raillard (1996)) molecules. Indeed, the development of in vitro selection and evolution techniques has made it possible to obtain novel ribozymes against a known substrate, using either random variants of a known ribozyme or random-sequence RNA as a starting point (Pan (1992); Tsang (1994); and Breaker (1994)).

Ribozymes, however, are highly susceptible to enzymatic hydrolysis within the cells where they are intended to perform their function. This in turn limits their pharmaceutical applications.

Recently, a new class of catalytic molecules called "DNAzymes" was created (Breaker and Joyce (1995); Santoro (1997)). DNAzymes are single-stranded, and cleave both RNA (Breaker (1994); Santoro (1997)) and DNA (Carmi (1996)). A general model for the DNAzyme has been proposed, and is known as the "10-23" model. DNAzymes following the "10-23" model, also

referred to simply as "10-23 DNazymes", have a catalytic domain of 15 deoxyribonucleotides, flanked by two substrate-recognition domains of seven to nine deoxyribonucleotides each. In vitro analyses show that this type of DNzyme can effectively cleave its substrate RNA at purine:pyrimidine
5 junctions under physiological conditions (Santoro (1997)).

DNazymes show promise as therapeutic agents. However, DNzyme success against a disease caused by the presence of a known mRNA molecule is not predictable. This unpredictability is due, in part, to two factors. First, certain mRNA secondary structures can impede a DNzyme's ability to bind
10 to and cleave its target mRNA. Second, the uptake of a DNzyme by cells expressing the target mRNA may not be efficient enough to permit therapeutically meaningful results. For these reasons, merely knowing of a disease and its causative target mRNA sequence does not alone allow one to reasonably predict the therapeutic success of a DNzyme against that target
15 mRNA, absent an inventive step.

SUMMARY OF THE INVENTION

Accordingly, in a first aspect the present invention provides a
20 DNzyme which specifically cleaves EGR-1 mRNA, the DNzyme including
(i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site;
(ii) a first binding domain contiguous with the 5' end of the catalytic domain; and
25 (iii) a second binding domain contiguous with the 3' end of the catalytic domain,

wherein the binding domains are sufficiently complementary to two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168 to 332 as shown in
30 SEQ ID NO:1, such that the DNzyme cleaves the EGR-1 mRNA.

In a second aspect the present invention provides a pharmaceutical composition including a DNzyme according to the first aspect and a pharmaceutically acceptable carrier.

In a third aspect the present invention provides a method of inhibiting
35 EGR-1 activity in cells which includes exposing the cells to a DNzyme according to the first aspect of the present invention.

WO 00/42173

PCT/AU00/00011

4

In a fourth aspect the present invention provides a method of inhibiting proliferation or migration of cells in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

5 In a fifth aspect the present invention provides a method of treating a condition associated with cell proliferation or migration in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

10 In a sixth aspect the present invention provides an angioplastic stent for inhibiting the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to the first aspect.

15 In a seventh aspect, the present invention provides a method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to the fifth aspect to the subject at around the time of the angioplasty.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 Sequence of NGFI-A DNzyme (ED5), its scrambled control (ED5SCR) and 23 nt synthetic rat substrate. The translational start site is underlined.

Figure 2 NGFI-A DNzyme inhibits the induction of NGFI-A mRNA and protein by serum. Northern blot analysis was performed with 25 μ g of total RNA. The blot was stripped and reprobed for β -Actin. Autoradiograms were analyzed by scanning densitometry and the ordinate axis is expressed as NGFI-A band intensity as a fraction of β -Actin band intensity. The mean and standard errors of the mean are indicated in the figure. Data is representative of 2 independent experiments. * indicates $P < 0.05$ (Student's paired t-test) as compared to control (FBS alone).

Figure 3 SMC proliferation is inhibited by NGFI-A DNzyme. **a**, Assessment of total cell numbers by Coulter counter. Growth-arrested SMCs that had been exposed to serum and/or DNzyme for 3 days were trypsinized followed by quantitation of the suspension. The sequence of AS2 is 5'-CTT GGC CGC TGC CAT-3' (SEQ ID NO: 20). **b**, Proportion of cells incorporating Trypan Blue after exposure to serum and/or DNzyme. Cells were stained incubated in 0.2% (w:v) Trypan Blue at 22 $^{\circ}$ C for 5 min prior to quantitation by hemocytometer in a blind manner. **c**, Effect of ED5 on pup SMC proliferation. Growth-arrested WKY12-22 cells exposed to serum and/or DNzyme for 3 days were resuspended and numbers were quantitated by Coulter counter. Data is representative of 2 independent experiments performed in triplicate. The mean and standard errors of the mean are indicated in the figure. * indicates $P < 0.05$ (Student's paired t-test) as compared to control (FBS alone).

Figure 4 NGFI-A DNzyme inhibition of neointima formation in the rat carotid artery. Neointimal and medial areas of 5 consecutive sections per rat (5 rats per group) taken at 250 μ m intervals from the point of ligation were determined digitally and expressed as a ratio per group. The mean and standard errors of the mean are indicated by the ordinate axis. * denotes

WO 00/42173

PCT/AU00/00011

6

P<0.05 as compared to the Lig, Lig+Veh or Lig+Veh+ED5SCR groups using the Wilcoxon rank sum test for unpaired data. Lig denotes ligation, Veh denotes vehicle.

5 **Figure 5** Selective inhibition of human smooth muscle cell proliferation by DzA.

Figure 6 Specific inhibition of porcine retinal smooth muscle cell proliferation by DzA.

10

In a preferred embodiment of the first aspect of the present invention,
35 the binding domains are complementary to the regions immediately flanking
the cleavage site. It will be appreciated by those skilled in the art, however,

- (i) 5'-caggggacaGGCTAGCTACAACGAacgttgcggg (SEQ ID NO: 3)
targets GU (nt 198, 199); arms hybridise to bp 189-207
- 35 (ii) 5'-tgcaggggaGGCTAGCTACAACGAaccgttgcg (SEQ ID NO: 4)
targets GU (nt 200, 201); arms hybridise to bp 191-209

- (iii) 5'-catcctggaGGCTAGCTACAACGAgagcaggct (SEQ ID NO: 5)
targets GU (nt 264, 265); arms hybridise to bp 255-273
- 5 (iv) 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6)
targets AU (nt 271, 272); arms hybridise to bp 262-280
- (v) 5'-ccgctgccaGGCTAGCTACAACGAcccgacgt (SEQ ID NO: 7)
targets AU (nt 271, 272); arms hybridise to bp 262-280
- 10 (vi) 5'-gcggggacaGGCTAGCTACAACGAcagctgcat (SEQ ID NO: 8)
targets AU (nt 301, 302); arms hybridise to bp 292-310
- (vii) 5'-cagcggggaGGCTAGCTACAACGAatcagctgc (SEQ ID NO: 9)
15 targets GU (nt 303, 304); arms hybridise to bp 294-312
- (viii) 5'-ggtcagagaGGCTAGCTACAACGActgcagcgg (SEQ ID NO: 10)
targets AU (nt 316, 317); arms hybridise to bp 307-325.

20 In a particularly preferred embodiment, the DNAzyme targets the AU site corresponding to nucleotides 271-272 (ie. the translation start codon).

In a further preferred embodiment, the DNAzyme has the sequence:
5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6).

In applying DNAzyme-based treatments, it is preferable that the
25 DNAzymes be as stable as possible against degradation in the intra-cellular milieu. One means of accomplishing this is by incorporating a 3'-3' inversion at one or more termini of the DNAzyme. More specifically, a 3'-3' inversion (also referred to herein simply as an "inversion") means the covalent phosphate bonding between the 3' carbons of the terminal nucleotide and its
30 adjacent nucleotide. This type of bonding is opposed to the normal phosphate bonding between the 3' and 5' carbons of adjacent nucleotides, hence the term "inversion". Accordingly, in a preferred embodiment, the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3' end of the catalytic domain. In addition to inversions, the instant
35 DNAzymes may contain modified nucleotides. Modified nucleotides

include, for example, N3'-P5' phosphoramidate linkages, and peptide-nucleic acid linkages. These are well known in the art.

In a particularly preferred embodiment, the DNAzyme includes an inverted T at the 3' position.

5 As will be appreciated by those skilled in the art, given that DNAzymes of the present invention cleave human EGR-1, similar DNAzymes can be produced to cleave the corresponding mRNA in other species, eg. rat (NGFI-A), mouse (Egr-1) etc. In a further aspect, the present invention provides such DNAzymes.

10 In a second aspect the present invention provides a pharmaceutical composition including a DNAzyme according to the first aspect and a pharmaceutically acceptable carrier.

In a third aspect the present invention provides a method of inhibiting EGR-1 activity in cells which includes exposing the cells to a DNAzyme
15 according to the first aspect of the present invention.

In a fourth aspect the present invention provides a method of inhibiting proliferation or migration of cells in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

20 In a fifth aspect the present invention provides a method of treating a condition associated with cell proliferation or migration in a subject which includes administering to the subject a prophylactically effective dose of a DNAzyme according to the first aspect of the present invention.

In preferred embodiments of the third, fourth and fifth aspects of the
25 present invention, the cells are vascular cells, particularly smooth muscle or endothelial cells. The cells may, however, be cells involved in neoplasia, such as tumour cells and the like.

Although the subject may be any animal or human, it is preferred that the subject is a human.

30 In a preferred embodiment, conditions associated with SMC proliferation (and migration) are selected from post-angioplasty restenosis, vein graft failure, transplant coronary disease and complications associated with atherosclerosis (cerebrovascular infarction (stroke), myocardial infarction (heart attack), hypertension or peripheral vascular disease
35 (gangrene of the extremities).

Within the parameters of the fourth and fifth aspects of the present invention, any suitable mode of administration may be used to administer or deliver the DNAzyme.

In particular, delivery of the nucleic acid agents described may be
5 achieved by one or more of the following methods:

- (a) Liposomes and liposome-protein conjugates and mixtures.
- (b) Using catheters to deliver intra-luminal formulations of the nucleic acid as a solution or in a complex with a liposome.
- (c) Catheter delivery to adventitial tissue as a solution or in a complex
10 with a liposome.
- (d) Within a polymer formulation such as polyethylenimine (PEI) or pluronic gels or within ethylene vinyl acetate copolymer (EVAc). The polymer is preferably delivered intra-luminally.
- (e) The nucleic acid may be bound to a delivery agent such as a
15 targetting moiety, or any suitable carrier such as a peptide or fatty acid molecule.
- (f) Within a viral-liposome complex, such as Sendai virus.
- (g) The nucleic acid may be delivered by a double angioplasty balloon device fixed to catheter.
- (h) The nucleic acid could be delivered on a specially prepared stent of
20 the Schatz-Palmaz or derivative type. The stent could be coated with a polymer or agent impregnated with nucleic acid that allows controlled release of the molecules at the vessel wall.

In a preferred embodiment, the mode of administration is topical
25 administration. Topical administration may be effected or performed using any of the various methods and delivery systems known to those skilled in the art. The topical administration can be performed, for example, via catheter and topical injection, and via coated stent as discussed below.

Pharmaceutical carriers for topical administration are well known in
30 the art, as are methods for combining same with active agents to be delivered. The following delivery systems, which employ a number of routinely used carriers, are only representative of the many embodiments envisioned for administering the instant composition.

Topical delivery systems include, for example, gels and solutions, and
35 can contain excipients such as solubilizers, permeation enhancers (e.g., fatty acids, fatty acid esters, fatty alcohols and amino acids), and hydrophilic

polymers (e.g., polycarbophil and polyvinylpyrrolidone). In the preferred embodiment, the pharmaceutically acceptable carrier is a liposome or a biodegradable polymer. Examples of agents which can be used in this invention include the following: (1) CellFectin, 1:1.5 (M/M) liposome formulation of the cationic lipid N,N^I,N^{II},N^{III}-tetramethyl-N,N^I,N^{II},N^{III}-tetrapalmitylspermine and dioleoyl phosphatidyl-ethanolamine (DOPE) (GIBCO BRL); (2) Cytofection GSV, 2:1 (M/M) liposome formulation of a cationic lipid and DOPE (Glen Research); (3) DOTAP (N-[1-(2,3-dioleoyloxy)-N,N,N-trimethyl-ammoniummethylsulfate) (Boehringer Mannheim); (4) Lipofectamine, 3:1 (M/M) liposome formulation of the polycationic lipid DOSPA and the neutral lipid DOPE (GIBCO BRL); (5) FuGENE[®] (Roche Molecular Biochemicals); (6) Superfect (Qiagen); and (7) Lipofectamine 2000 (Gibco-life Technologies).

Examples of suitable methods for topical administration of the DNAzymes of the present invention are described in Autieri et al. (1995), Simons et al. (1992), Morishita et al. (1993), Bennett and Schwartz (1995) and Frimerman et al. (1999).

Determining the prophylactically effective dose of the instant pharmaceutical composition can be done based on animal data using routine computational methods. In one embodiment, the prophylactically effective dose contains between about 0.1 mg and about 1 g of the instant DNAzyme. In another embodiment, the prophylactically effective dose contains between about 1 mg and about 100 mg of the instant DNAzyme. In a further embodiment, the prophylactically effective dose contains between about 10 mg and about 50 mg of the instant DNAzyme. In yet a further embodiment, the prophylactically effective does contains about 25 mg of the instant DNAzyme.

In a sixth aspect the present invention provides an angioplastic stent for inhibiting the onset of restenosis, which comprises an angioplastic stent operably coated with a prophylactically effective dose of a DNAzyme according to the first aspect.

Angioplastic stents, also known by other terms such as "intravascular stents" or simple "stents", are well known in the art. They are routinely used to prevent vascular closure due to physical anomalies such as unwanted inward growth of vascular tissue due to surgical trauma. They often have a

tubular, expanding lattice-type structure appropriate for their function, and can optionally be biodegradable.

In this invention, the stent can be operably coated with the instant pharmaceutical composition using any suitable means known in the art.

5 Here, "operably coating" a stent means coating it in a way that permits the timely release of the pharmaceutical composition into the surrounding tissue to be treated once the coated stent is administered. Such coating methods, for example, can use the polymer polypyrrole.

10 In a seventh aspect, the present invention provides a method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to the fifth aspect to the subject at around the time of the angioplasty.

As used herein, administration "at around the time of angioplasty" can be performed during the procedure, or immediately before or after the procedure. The administering can be performed according to known 15 methods such as catheter delivery.

In order that the nature of the present invention may be more clearly understood, preferred forms thereof will now be described with reference to the following non-limiting Figures and Examples.

```

5      Symbol comparison table: GenRunData:pileupdna.cmp  CompCheck: 6876
          GapWeight: 5.000
          GapLengthWeight: 0.300
      EGR1align.msf  MSF: 4388  Type: N  April  7, 1998 12:07  Check: 5107

10     Name: mouseEGR1  Len:  4388  Check: 8340  Weight: 1.0 (SEQ ID NO:11)
      Name: ratEGR1    Len:  4388  Check: 8587  Weight: 1.0 (SEQ ID NO:12)
      Name: humanEGR1  Len:  4388  Check: 8180  Weight: 1.00 (SEQ ID NO:1)

      NB. THIS IS RAT NGFI-A numbering

15     1 50
      mouseEgr1 .....
      ratNGFIA CCGCGGAGCC TCAGCTCTAC GCGCCTGGCG CCCTCCCTAC GCGGGCGTCC
      humanEGR1 .....

20     51 100
      mouseEGR1 .....
      ratEGR1  CCGACTCCCG CGCGCGTTCA GGCTCCGGGT TGGGAACCAA GGAGGGGGAG
      humanEGR1 .....

25     101 150
      mouseEGR1 .....
      ratEGR1  GGTGGGTGCG CCGACCCGGA AACACCATAT AAGGAGCAGG AAGGATCCCC
      humanEGR1 .....

30     151 200
      mouseEGR1 .....
      ratEGR1  CGCCGGAACA GACCTTATTT GGGCAGCGCC TTATATGGAG TGGCCCAATA
      humanEGR1 .....

35     201 250
      mouseEGR1 .....
      ratEGR1  TGGCCCTGCC GCTTCCGGCT CTGGGAGGAG GGGCGAACGG GGGTTGGGGC
      humanEGR1 .....

40     251 300
      mouseEGR1 .....
      ratEGR1  GGGGGCAAGC TGGGAACTCC AGGAGCCTAG CCCGGGAGGC CACTGCCGCT
      humanEGR1 .....

45     301 350
      mouseEGR1 .....
      ratEGR1  GTTCCAATAC TAGGCTTTCC AGGAGCCTGA GCGCTCAGGG TGCCGGAGCC
      humanEGR1 .....

50     351 400
      mouseEGR1 .....
      ratEGR1  GGTCGCAGGG TGGAAGCGCC CACCGCTCTT GGATGGGAGG TCTTCACGTC
      humanEGR1 .....

55     401 450
      mouseEGR1 .....
      ratEGR1  ACTCCGGGTC CTCCCGGTCTG GTCCTTCCAT ATTAGGGCTT CCTGCTTCCC
      humanEGR1 .....

60     451 500

```

WO 00/42173

PCT/AU00/00011

15

	mouseEGR1
	ratEGR1	ATATATGGCC	ATGTACGTCA	CGGCGGAGGC	GGGCCCCTGC	TGTTTCAGAC
	humanEGR1
5		501				550
	mouseEGR1
	ratEGR1	CCTTGAAATA	GAGGCCGATT	CGGGGAGTCG	CGAGAGATCC	CAGCGCGCAG
	humanEGR1CCGCAG
10		551				600
	mouseEGR1GGGGA	GCCGCCGCCG	CGATTTCGCCG	CCGCCGCCAG	CTTCCGCCGC
	ratEGR1	AACTTGGGGA	GCCGCCGCCG	CGATTTCGCCG	CCGCCGCCAG	CTTCCGCCGC
	humanEGR1	AACTTGGGGA	GCCGCCGCCG	CCATCCGCCG	CCGCAGCCAG	CTTCCGCCGC
15		601				650
	mouseEGR1	CGCAAGATCG	GCCCCTGCCC	CAGCCTCCGC	GGCAGCCCTG	CGTCCACCAC
	ratEGR1	CGCAAGATCG	GCCCCTGCCC	CAGCCTCCGC	GGCAGCCCTG	CGTCCACCAC
	humanEGR1	CGCAGGACCG	GCCCCTGCCC	CAGCCTCCGC	AGCCGCGGCG	CGTCCACGCC
20		651				700
	mouseEGR1	GGGCCGCGGC	TACCGCCAGC	CTGGGGGGCCC	ACCTACACTC	CCCGCAGTGT
	ratEGR1	GGGCCGCGGC	CACCGCCAGC	CTGGGGGGCCC	ACCTACACTC	CCCGCAGTGT
	humanEGR1	CGCCCGCGCC	CAGGGCGAGT	CGGGGTCGCC	GCCTGCACGC	TTCTCAGTGT
25		701				750
	mouseEGR1	GCCCCTGCAC	CCCGCATGTA	ACCCGGCCAA	CCCCCGGCGA	GTGTGCCCTC
	ratEGR1	GCCCCTGCAC	CCCGCATGTA	ACCCGGCCAA	CATCCGGCGA	GTGTGCCCTC
	humanEGR1	TCCCC.GCGC	CCCGCATGTA	ACCCGGCCAG	GCCCCCGCAA	CGGTGTCCCC
30		751				800
	mouseEGR1	AGTAGCTTCG	GCCCCGGGCT	GCGCCCACC.	.ACCCAACAT	CAGTTCTCCA
	ratEGR1	AGTAGCTTCG	GCCCCGGGCT	GCGCCCACC.	.ACCCAACAT	CAGCTCTCCA
	humanEGR1	TGCAGCTCCA	GCCCCGGGCT	GCACCCCCC	GCCCCGACAC	CAGCTCTCCA
35		801				850
	mouseEGR1	GCTCGCTGGT	CCGGGATGGC	AGCGGCCAAG	GCCGAGATGC	AATTGATGTC
	ratEGR1	GCTCGCACGT	CCGGGATGGC	AGCGGCCAAG	GCCGAGATGC	AATTGATGTC
	humanEGR1	GCCTGCTCGT	CCAGGATGGC	CGCGGCCAAG	GCCGAGATGC	AGCTGATGTC
40		ED5 (rat) arms hybridise to bp 807-825 in rat sequ hED5(hum) arms hybridise to bp 262-280 in hum sequ				
		851				900
	mouseEGR1	TCCGCTGCAG	ATCTCTGACC	CGTTCGGGCTC	CTTTCCTCAC	TCACCCACCA
45	ratEGR1	TCCGCTGCAG	ATCTCTGACC	CGTTCGGGCTC	CTTTCCTCAC	TCACCCACCA
	humanEGR1	CCCGCTGCAG	ATCTCTGACC	CGTTCGGATC	CTTTCCTCAC	TCGCCCACCA
		901				950
	mouseEGR1	TGGACAACATA	CCCCAAACTG	GAGGAGATGA	TGCTGCTGAG	CAACGGGGCT
50	ratEGR1	TGGACAACATA	CCCCAAACTG	GAGGAGATGA	TGCTGCTGAG	CAACGGGGCT
	humanEGR1	TGGACAACATA	CCCTAAGCTG	GAGGAGATGA	TGCTGCTGAG	CAACGGGGCT
		951				1000
	mouseEGR1	CCCCAGTTCC	TCGGTGCTGC	CGGAACCCCA	GAGGGCAGCG	GCGGTAAT..
55	ratEGR1	CCCCAGTTCC	TCGGTGCTGC	CGGAACCCCA	GAGGGCAGCG	GCGGCAATAA
	humanEGR1	CCCCAGTTCC	TCGGCGCCGC	CGGGGCCCA	GAGGGCAGCG	GCAGCAACAG
		1001				1050
	mouseEGR1AGC	AGCAGCAGCA	CCAGCAGCGG	GGGCGGTGGT	GGGGGCGGCA
60	ratEGR1	CAGCAGCAGC	AGCAGCAGCA	GCAGCAGCGG	GGGCGGTGGT	GGGGGCGGCA

WO 00/42173

PCT/AU00/00011

16

	humanEGR1	CAGCAGCAGC	AGCAGCGGGG	GC	CGGTGGAGG	CGGCGGGGGC	GGCAGCAACA
		1051					1100
5	mouseEGR1	GCAACAGCGG	CAGCAGCGCC	TTCAATCCTC	AAGGGGAGCC	GAGCGAACAA	
	rateEGR1	GCAACAGCGG	CAGCAGCGCT	TTCAATCCTC	AAGGGGAGCC	GAGCGAACAA	
	humanEGR1	GCAGCAGCAG	CAGCAGCACC	TTCAACCCTC	AGGCGGACAC	GGGCGAGCAG	
		1101					1150
10	mouseEGR1	CCCTATGAGC	ACCTGACCAC	AG...AGTCC	TTTTCTGACA	TCGCTCTGAA	
	rateEGR1	CCCTACGAGC	ACCTGACCAC	AGGTAAGCGG	TGGTCTGCGC	CGAGGCTGAA	
	humanEGR1	CCCTACGAGC	ACCTGACCGC	AG...AGTCT	TTTCCTGACA	TCTCTCTGAA	
		1151					1200
15	mouseEGR1	TAATGAGAAG	GCGATGGTGG	AGACGAGTTA	TCCCAGCCAA	ACGACTCGGT	
	rateEGR1	TCCCCCTTCG	TGACTACCTT	AACGTCCAGT	CCTTTGCAGC	ACGGACCTGC	
	humanEGR1	CAACGAGAAG	GTGCTGGTGG	AGACCAGTTA	CCCCAGCCAA	ACCACTCGAC	
		1201					1250
20	mouseEGR1	TGCCTCCCAT	CACCTATACT	GGCCGCTTCT	CCCTGGAGCC	CGCACCCAAC	
	rateEGR1	ATCTAGATCT	TAGGGACGGG	ATTGGGATTT	CCCTCTATTC	..CACACAGC	
	humanEGR1	TGCCCCCAT	CACCTATACT	GGCCGCTTTT	CCCTGGAGCC	TGCACCCAAC	
		1251					1300
25	mouseEGR1	AGTGGCAACA	CTTTGTGGCC	TGAACCCCTT	TTCAGCCTAG	TCAGTGGCCT	
	rateEGR1	TCCAGGGACT	TGTGTTAGAG	GGATGTCTGG	GGACCCCCCA	ACCCTCCATC	
	humanEGR1	AGTGGCAACA	CCTTGTGGCC	CGAGCCCCCT	TTCAGCTTGG	TCAGTGGCCT	
		1301					1350
30	mouseEGR1	CGTGAGCATG	ACCAATCCTC	CGACCTCTTC	ATCCTCGGCG	CCTTCTCCAG	
	rateEGR1	CTTGCGGGTG	CGCGGAGGGC	AGACCGTTTG	TTTTGGATGG	AGAACTCAAG	
	humanEGR1	AGTGAGCATG	ACCAACCCAC	CGGCCTCCTC	GTCCTCAGCA	CCATCTCCAG	
		1351					1400
35	mouseEGR1	CTGCTTCATC	GTCTTCCTCT	GCCTCCCAGA	GCCCGCCCCCT	GAGCTGTGCC	
	rateEGR1	TTGCGTGGGT	GGCT.....GGAGT	GGGGGAGGGT	TTGTTTTGAT	
	humanEGR1	CGGCCTCCTC	CGC...CTCC	GCCTCCCAGA	GCCCACCCCT	GAGCTGCGCA	
		1401					1450
40	mouseEGR1	GTGCCGTCCA	ACGACAGCAG	TCCCATCTAC	TCGGCTGCGC	CCACCTTTCC	
	rateEGR1	GAGCAGGGTT	GC....CCCC	TCCCCCGCGC	GCGTTGTGCG	GAGCCTTGTT	
	humanEGR1	GTGCCATCCA	ACGACAGCAG	TCCCATTTAC	TCAGCGGCAC	CCACCTTCCC	
		1451					1500
45	mouseEGR1	TACTCCCAAC	ACTGACATTT	TTCCTGAGCC	CCAAAGCCAG	GCCTTTCCCTG	
	rateEGR1	TGCAGCTTGT	TCCCAAGGAA	GGGCTGAAAT	CTGTCACCAG	GGATGTCCCG	
	humanEGR1	CACGCCGAAC	ACTGACATTT	TCCCTGAGCC	ACAAAGCCAG	GCCTTCCCCG	
		1501					1550
50	mouseEGR1	GCTCGGCAGG	CACAGCCTTG	CAGTACCCGC	CTCCTGCCTA	CCCTGCCACC	
	rateEGR1	CCGCCCAGGG	TAGGGGCGCG	CATTAGCTGT	GGCC.ACTAG	GGTGCTGGCG	
	humanEGR1	GCTCGGCAGG	GACAGCGCTC	CAGTACCCGC	CTCCTGCCTA	CCCTGCCGCC	
		1551					1600
55	mouseEGR1	AAAGGTGGTT	TCCAGGTTCC	CATGATCCCT	GACTATCTGT	TTCCACAACA	
	rateEGR1	GGATTCCCTC	ACCCCGGACG	CCTGCTGCGG	AGCGCTCTCA	GAGCTGCAGT	
	humanEGR1	AAGGGTGGCT	TCCAGGTTCC	CATGATCCCC	GACTACCTGT	TTCCACAGCA	
		1601					1650
60	mouseEGR1	ACAGGGAGAC	CTGAGCCTGG	GCACCCCAAG	CCAGAAGCCC	TTCCAGGGTC	
	rateEGR1	AGAGGGGGAT	TCTCTGTTTG	CGTCAGCTGT	CGAAATGGCT	CT.....GC	

	humanEGR1	GCAGGGGGGAT	CTGGGGCCTGG	GCACCCCAGA	CCAGAAGCCC	TTCCAGGGCC
		1651				1700
5	mouseEGR1	TGGAGAACCG	TACCCAGCAG	CCTTCGCTCA	CTCCACTATC	CACTATTAATA
	rateEGR1	CACTGGAGCA	GGTCCAGGAA	CATTGCAATC	TGCTGCTATC	AATTATTAAC
	humanEGR1	TGGAGAGCCG	CACCCAGCAG	CCTTCGCTAA	CCCCTCTGTC	TACTATTAAG
		1701				1750
10	mouseEGR1	GCCTTCGCCA	CTCAGTCGGG	CTCCCAGGAC	TTAAAG....	...GCTCTTA
	rateEGR1	CACATCGAGA	GTCAGTGGTA	GCCGGGCGAC	CTCTTGCCCTG	GCCGCTTCGG
	humanEGR1	GCCTTTGCCA	CTCAGTCGGG	CTCCCAGGAC	CTGAAG....	...GCCCTCA
		1751				1800
15	mouseEGR1	ATACCACCTA	CCAATCCCAG	CTCATCA..A	ACCCAGCCGC	ATGCGCAAGT
	rateEGR1	CTCTCATCGT	CCAGTGATTG	CTCTCCAGTA	ACCAGGCCCTC	TCTGTTCTCT
	humanEGR1	ATACCAGCTA	CCAGTCCCAG	CTCATCA..A	ACCCAGCCGC	ATGCGCAAGT
		1801				1850
20	mouseEGR1	ACCCCAACCG	GCCCAGCAAG	ACACCCCCCC	ATGAACGCCC	ATATGCTTGC
	rateEGR1	TTCCTGCCAG	AGTCCTTTTC	TGACATCGCT	CTGAATAACG	AGAAG..GCG
	humanEGR1	ATCCCAACCG	GCCCAGCAAG	ACGCCCCCCC	ACGAACGCCC	TTACGCTTGC
		1851				1900
25	mouseEGR1	CCTGTCGAGT	CCTGCGATCG	CCGCTTTTCT	CGCTCGGATG	AGCTTACCCG
	rateEGR1	CTGGTGGAGA	CAAGTTATCC	CAGCCAAACT	ACCCGGTTGC	CTCCCATCAC
	humanEGR1	CCAGTGGAGT	CCTGTGATCG	CCGCTTCTCC	CGCTCCGACG	AGCTCACCCG
		1901				1950
30	mouseEGR1	CCATATCCGC	ATCCACACAG	GCCAGAAGCC	CTTCCAGTGT	CGAATCTGCA
	rateEGR1	CTATACTGGC	CGCTTCTCCC	TGGAGCCTGC	ACCCAACAGT	GGCAACACTT
	humanEGR1	CCACATCCGC	ATCCACACAG	GCCAGAAGCC	CTTCCAGTGC	CGCATCTGCA
		1951				2000
35	mouseEGR1	TGCGTAACTT	CAGTCGTAGT	GACCACCTTA	CCACCCACAT	CCGCACCCAC
	rateEGR1	TGTGGCCTGA	ACCCCTTTTC	AGCCTAGTCA	GTGGCCTTGT	GAGCATGACC
	humanEGR1	TGCGCAACTT	CAGCCGCAGC	GACCACCTCA	CCACCCACAT	CCGCACCCAC
		2001				2050
40	mouseEGR1	ACAGGCGAGA	AGCCTTTTGC	CTGTGACATT	TGTGGGAGGA	AGTTTGCCAG
	rateEGR1	AACCCTCCAA	CCTCTTCATC	CTCAGCGCCT	TCTCCAGCTG	CTTCATCGTC
	humanEGR1	ACAGGCGAAA	AGCCCTTCGC	CTGCGACATC	TGTGGAAGAA	AGTTTGCCAG
		2051				2100
45	mouseEGR1	GAGTGATGAA	CGCAAGAGGC	ATACCAAAAT	CCATTTAAGA	CAGAAGGACA
	rateEGR1	TTCTCTTGCC	TCCCAGAGCC	CACCCCTGAG	CTGTGCCGTG	CCGTCCAACG
	humanEGR1	GAGCGATGAA	CGCAAGAGGC	ATACCAAGAT	CCACTTGCGG	CAGAAGGACA
		2101				2150
50	mouseEGR1	AGAAAGCAGA	CAAAAGTGTG	GTGGCCTCCC	CGGCTGC...	.CTCTTCACT
	rateEGR1	ACAGCAGTCC	CATTTACTCA	GCTGCACCCA	CCTTTTCTAC	TCCCAACACT
	humanEGR1	AGAAAGCAGA	CAAAAGTGTT	GTGGCCTCTT	CGGCCACCTC	CTCTCTCTCT
		2151				2200
55	mouseEGR1	CTCTTCTTAC	CCATCCCCAG	TGGCTACCTC
	rateEGR1	GACATTTTTC	CTGAGCCCCA	AAGCCAGGCC
	humanEGR1	TCCTACCCGT	CCCCGGTTGC	TACCTCTTAC	CCGTCCCCGG	TTACTACCTC
		2201				2250
60	mouseEGR1	CTACCCATCC	CCTGCCACCA	CCTCATTTCC	ATCCCCTGTG	CCCCTTCTCT
	rateEGR1	TTTCCTGGCT	CTGCAGGCAC	AGCCTTGCAG	TACCCGCTTC	CTGCCTACCC
	humanEGR1	CTACCCATCC	CCTGCCACCA	CCTCATTTCC	ATCCCCTGTG	CCCCTTCTCT

WO 00/42173

PCT/AU00/00011

18

	humanEGR1	TTATCCATCC	CCGGCCACCA	CCTCATACCC	ATCCCCTGTG	CCCACCTCCT
		2251				2300
5	mouseEGR1	ACTCCTCTCC	TGGCTCCTCC	ACCTACCCAT	CTCCTGCGCA	CAGTGGCTTC
	ratEGR1	TGCCACCAAG	GGTGGTTTCC	AGGTTCCTAT	GATCCCTGAC	TATCTGTTTC
	humanEGR1	TCTCCTCTCC	CGGCTCCTCG	ACCTACCCAT	CCCCTGTGCA	CAGTGGCTTC
		2301				2350
10	mouseEGR1	CCGTCGCCGT	CAGTGGCCAC	CACCTTTGCC	TCCGTTCC..
	ratEGR1	CACAACAACA	GGGAGACCTG	AGCCTGGGCA	CCCCAGACCA	GAAGCCCTTC
	humanEGR1	CCCTCCCCGT	CGGTGGCCAC	CACGTACTCC	TCTGTTCCC.
		2351				2400
15	mouseEGR1ACCTGC	TTTCCCCACC	CAGGTCAGCA	GCTTCCCGTC	TGCGGGCGTC
	ratEGR1	CAGGGTCTGG	AGAACCGTAC	CCAGCAGCCT	TCGCTCACTC	CACTATCCAC
	humanEGR1CCTGC	TTTCCCGGCC	CAGGTCAGCA	GCTTCCCTTC	CTCAGCTGTC
		2401				2450
20	mouseEGR1	AGCAGCTCCT	TCAGCACCTC	AACTGGTCTT	TCAGACATGA	CAGCGACCTT
	ratEGR1	TATCAAAGCC	TTCGCCACTC	AGTCGGGCTC	CCAGGACTTA	AAGGCTCTTA
	humanEGR1	ACCAACTCCT	TCAGCGCCTC	CACAGGGCTT	TCGGACATGA	CAGCAACCTT
		2451				2500
25	mouseEGR1	TTCTCCCAGG	ACAATTGAAA	TTTGCTAAAG	GGA.....	.ATAAAAG..
	ratEGR1	ATAACACCTA	CCAGTCCCAA	CTCATCAAAC	CCAGCCGCAT	GCGCAAGT..
	humanEGR1	TTCTCCCAGG	ACAATTGAAA	TTTGCTAAAG	GGAAAGGGGA	AAGAAAGGGA
		2501				2550
30	mouseEGR1	.AAAGCAAAG	GGAGAGGCAG	GAAAGACATA	AAAGCA...C	AGGAGGGAAG
	ratEGR1	.ACCCCAAAC	GGCCACAGCA	GACACCCCCC	CATGAACGCC	CGTATGCTTG
	humanEGR1	AAAGGGAGAA	AAAGAAACAC	AAGAGACTTA	AAGGACAGGA	GGAGGAGATG
		2551				2600
35	mouseEGR1	AGATGGCCGC	AAGAGGGGCC	ACCTCTTAGG	TCAGATGGAA	GATCTCAGAG
	ratEGR1	CCCTGTTGAG	TCCTGCGATC	GCCGCTTTTC	TCGCTCGGAT	GAGCTTACAC
	humanEGR1	GCCATAGGAG	AGGAGGGTT.	.CCTCTTAGG	TCAGATGGAG	GTTCTCAGAG
		2601				2650
40	mouseEGR1	CCAAGTCCTT	CTACTCACGA	GTA..GAAGG	ACCGTTGGCC	AACAGCCCTT
	ratEGR1	GCCACATCCG	CATCCATACA	GGC..CAGAA	GCCCTTCCAG	TGTCGAATCT
	humanEGR1	CCAAGTCCTC	CCTCTCTACT	GGAGTGGAAG	GTCTATTGGC	CAACAATCCT
		2651				2700
45	mouseEGR1	TCACTTACCA	TCCCTGCCTC	CCCCGTCTTG	TTCCCTTTGA	CTTCAGCTGC
	ratEGR1	GCAATGCGTA	TTTCAGTCGT	AGTGACCACC	TTACCACCCA	CATCCGCACC
	humanEGR1	TTCTGCCCCAC	TTCCCTTTCC	CCAATTACTA	TTCCCTTTGA	CTTCAGCTGC
		2701				2750
50	mouseEGR1	CTGAAACAGC	CATGTCCAAG	TTCTTCACCT	CTATCCAAAG	GAATTGATTT
	ratEGR1	C..ACACAGG	CGAGAAGCCT	TTTGCCCTGTG	ACATTTGTGG	GAGAAAGTTT
	humanEGR1	CTGAAACAGC	CATGTCCAAG	TTCTTCACCT	CTATCCAAAG	AACTTGATTT
		2751				2800
55	mouseEGR1	GCAATG....	..TATTGGAT	AAATCATTTT	AGTATCCTCT
	ratEGR1	GCCAGGAGTG	ATGAACGCAA	GAGGCATACC	AAAATCCACT	TAAGACAGAA
	humanEGR1	GCAATGGA...	..TTTTGGAT	AAATCATTTT	AGTATCATCT
		2801				2850
60	mouseEGR1CCATC	ACATGCCTGG	CCCTTGCTCC	CTTCAGCGCT	AGACCATCAA
	ratEGR1	GGACAAGAAA	GCAGACAAAA	GTGTCGTGGC	CTCCTCAGCT	GCCTCTTCCC

WO 00/42173

PCT/AU00/00011

19

	humanEGR1CCATCA	TATGCCTGAC	CCCTTGCTCC	CTTCAATGCT	AGAAAATCGA
		2851				2900
5	mouseEGR1	GTTGGCATAA	AGAAAAAAA	ATGGGTTTGG	GCCCTCAGAA	CCCTGCCCTG
	ratEGR1	TCTCTTCCTA	CCCATCCCCA	GTGGCTACCT	CCTACCCATC	CCCCGCCACC
	humanEGR1	GTTGGC....AAAAT	GGGGTTTGGG	CCCCTCAGAG	CCCTGCCCTG
		2901				2950
10	mouseEGR1	CATCTTTGTA	CAGCATCTGT	GCCATGGATT	TTGTTTTTCCT	TGGGGTATTC
	ratEGR1	ACCTCATTTT	CATCCCCAGT	GCCCACCTCT	TACTCCTCTC	CGGGCTCCTC
	humanEGR1	CACCCTTGTA	CAGTGTCTGT	GCCATGGATT	TCGTTTTTCT	TGGGGTACTC
		2951				3000
15	mouseEGR1	TTGATGTGAA	GATAATTTGC	ATACT.....	.CTATTGTAT	TATTTGGAGT
	ratEGR1	TACCTACCCG	TCTCCTGCAC	ACAGTGGCTT	CCCATCGCCC	TCGGTGGCCA
	humanEGR1	TTGATGTGAA	GATAATTTGC	ATATT.....	.CTATTGTAT	TATTTGGAGT
		3001				3050
20	mouseEGR1	TAAATCCTCA	CTTTGGGG..	GAGGGGGGAG	CAAAGCCAAG	CAAACCAATG
	ratEGR1	CCACCTATGC	CTCCGTCC..	CACCTGCTTT	CCCTGCCCAG	GTCAGCACCT
	humanEGR1	TAGGTCCTCA	CTTGGGGGAA	AAAAAAAAAA	AAAAGCCAAG	CAAACCAATG
		3051				3100
25	mouseEGR1	ATGATCCTCT	ATTTTGTGAT	GACTCTGCTG	TGACATTA..
	ratEGR1	TCCAGTCTGC	AGGGGTGAGC	AACTCCTTCA	GCACCTCAAC	GGGTCTTTCA
	humanEGR1	GTGATCCTCT	ATTTTGTGAT	GATGCTGTGA	CAATA.....
		3101				3150
30	mouseEGR1	.GGTTTGAAG	CATTTTTTTT	TTCAAGCAGC	AGTCCTAGGT	ATTAAGTGA
	ratEGR1	GACATGACAG	CAACCTTTTC	TCCTAGGACA	ATTGAAATTT	GCTAAAGGGA
	humanEGR1	...AGTTTGA	ACCTTTTTTT	TTGAAACAGC	AGTCCCAG..	..TATTCTCA
		3151				3200
35	mouseEGR1	..GCATGTGT	CAGAGTGTTG	TTCCGTTAAT	TTTGTAATA	CTGGCTCGAC
	ratEGR1	ATGAAAGAGA	GCAAAGGGAG	GGGAGCGCGA	GAGACAATAA	AGGACAGGAG
	humanEGR1	GAGCATGTGT	CAGAGTGTTG	TTCCGTTAAC	CTTTTTGTAA	ATACTGCTTG
		3201				3250
40	mouseEGR1	.TGTAACCTCT	CACATGTGAC	AAAGTATGGT	TTGTTTGGTT	GGGTTTTGTT
	ratEGR1	.GGAAGAAAT	GGCCCGCAAG	AGGGGCTGCC	TCTTAGGTCA	GATGGAAGAT
	humanEGR1	ACCGTACTCT	CACATGTGGC	AAAATATGGT	TTGGTTTTTC	TTTTTTTTTT
		3251				3300
45	mouseEGR1	TTTGAGAATT	TTTTTGCCCG	TCCCTTTGGT	TTCAAAAGTT	TCACGTCTTG
	ratEGR1	CTCAGAGCCA	AGTCCTTCTA	GTCAGTAGAA	GGCCCGTTGG	CCACCAGCCC
	humanEGR1	TTGAAAGTGT	TTTTTCTTCG	TCCTTTTGGT	TTAAAAAGTT	TCACGTCTTG
		3301				3350
50	mouseEGR1	GTGCCTTTTG	TGTGACACGC	CTT.CCGATG	GCTTGACATG	CGCA.....
	ratEGR1	TTTCACTTAG	CGTCCCTGCC	CTC.CCCAGT	CCCGGTCCCT	TTGACTTCAG
	humanEGR1	GTGCCTTTTG	TGTGATGCCC	CTTGCTGATG	GCTTGACATG	TGCAAT....
		3351				3400
55	mouseEGR1	...GATGTGA	GGGACACGCT	CACCTTAGCC	TTAA...GGG	GGTAGGAGTG
	ratEGR1	CTGCCTGAAA	CAGCCACGTC	CAAGTTCTTC	ACCT...CTA	TCCAAAGGAC
	humanEGR1TGTGA	GGGACATGCT	CACCTCTAGC	CTTAAGGGGG	GCAGGGAGTG
		3401				3450
60	mouseEGR1	ATGTGTTGGG	GGAGGCTTGA	GAGCAAAAC	GAGGAAGAGG	GCTGAGCTGA
	ratEGR1	TTGATTTGCA	TGGTATTGGA	TAAACCATTT	CAGCATCATC	TCCACCACAT

WO 00/42173

PCT/AU00/00011

20

	humanEGR1	ATGATTTGGG	GGAGGCTTTG	GGAGCAAAAT	AAGGAAGAGG	GCTGAGCTGA	
		3451				3500	
5	mouseEGR1	GCTTTCGGTC	TCCAGAATGT	AAGAAGAAAA	AATTTAAACA	AAAATCTGAA	
	ratEGR1	GCCTGGCCCT	TGCTCCCTTC	AGCACTAGAA	CATCAAGTTG	GCTGAAAAAA	
	humanEGR1	GCTTCGGTTC	TCCAGAATGT	AAGAAAACAA	AATCTAAAAC	AAAATCTGAA	
		3501				3550	
10	mouseEGR1	CTCTCAAAAG	TCTATTTTTC	TAAACTGAAA	ATGTAAATTT	ATACATCTAT	
	ratEGR1	AAAATGGGTC	TGGGCCCTCA	GAACCCTGCC	CTGTATCTTT	GTACA.....	
	humanEGR1	CTCTCAAAAG	TCTATTTTTC	TAA.CTGAAA	ATGTAAATTT	ATAAATATAT	
		3551				3600	
15	mouseEGR1	TCAGGAGTTG	GAGTGTTGTG	GTTACCTACT	GAGTAGGCTG	CAGTTTTTGT	
	ratEGR1	GCATCTGTGC	CATGGATTTT	GTTTTCTTGT	GGGTATTCTT	GATGTGAAGA	
	humanEGR1	TCAGGAGTTG	GAATGTTGTA	GTTACCTACT	GAGTAGGCGG	CGATTTTTGT	
		3601				3650	
20	mouseEGR1	ATGTTATGAA	CATGAAGTTC	ATTATTTTGT	GGTTTTATTT	TACTTTGTAC	
	ratEGR1	TAATTTGCAT	ACTCTATTGT	ACTATTTGGA	GTAAATTCTT	CACTTTGGGG	
	humanEGR1	ATGTTATGAA	CATGCAGTTC	ATTATTTTGT	GGTCTATTTT	TACTTTGTAC	
		3651				3700	
25	mouseEGR1	TTGTGTTTGC	TTAAACAAAG	TAACCTGTTT	GGCTTATAAA	CACATTGAAT	
	ratEGR1	GAGGGGGAGC	AAAGCCAAGC	AAACCAATGG	TGATCCTCTA	TTTTGTGATG	
	humanEGR1	TTGTGTTTGC	TTAAACAAAG	TGA.CTGTTT	GGCTTATAAA	CACATTGAAT	
		3701				3750	
30	mouseEGR1	GCGCTCTATT	GCCCATGG..	..GATATGTG	GTGTGTATCC	TTCAGAAAAA	
	ratEGR1	ATCCTGCTGT	GACATTAGGT	TTGAAACTTT	TTTTTTTTTT	TGAAGCAGCA	
	humanEGR1	GCGCTTTATT	GCCCATGG..	..GATATGTG	GTGTATATCC	TTCCAAAAAA	
		3751				3800	
35	mouseEGR1	TTAAAAGGAA	AAAT.....	
	ratEGR1	GTCCTAGGTA	TTAACTGGAG	CATGTGTCAG	AGTGTGTGTC	CGTTAATTTT	
	humanEGR1	TTAAAACGAA	AATAAAGTAG	CTGCGATTGG	G.....	
		3801				3850	
40	mouseEGR1	
	ratEGR1	GTAAATACTG	CTCGACTGTA	ACTCTCACAT	GTGACAAAAT	ACGGTTTGTG	
	humanEGR1	
		3851				3900	
45	mouseEGR1	
	ratEGR1	TGGTTGGGTT	TTTTGTGTGT	TTTGAAAAAA	AAATTTTTTT	TTTGCCCGTC	
	humanEGR1	
		3901				3950	
50	mouseEGR1	
	ratEGR1	CCTTTGGTTT	CAAAAGTTTC	ACGTCTTGGT	GCCTTTGTGT	GACACACCTT	
	humanEGR1	
		3951				4000	
55	mouseEGR1	
	ratEGR1	GCCGATGGCT	GGACATGTGC	AATCGTGAGG	GGACACGCTC	ACCTCTAGCC	
	humanEGR1	
		4001				4050	
60	mouseEGR1	
	ratEGR1	TTAAGGGGGT	AGGAGTGATG	TTTCAGGGGA	GGCTTTAGAG	CACGATGAGG	

WO 00/42173

PCT/AJ00/00011

21

	humanEGR1
		4051				4100
5	mouseEGR1
	ratEGR1	AAGAGGGCTG	AGCTGAGCTT	TGGTTCTCCA	GAATGTAAGA	AGAAAAATTT
	humanEGR1
		4101				4150
10	mouseEGR1
	ratEGR1	AAAACAAAAA	TCTGAACTCT	CAAAAGTCTA	TTTTTTTAAC	TGAAAATGTA
	humanEGR1
		4151				4200
15	mouseEGR1
	ratEGR1	GATTTATCCA	TGTTCGGGAG	TTGGAATGCT	GCGGTTACCT	ACTGAGTAGG
	humanEGR1
		4201				4250
20	mouseEGR1
	ratEGR1	CGGTGACTTT	TGTATGCTAT	GAACATGAAG	TTCATTATTT	TGTGGTTTTA
	humanEGR1
		4251				4300
25	mouseEGR1
	ratEGR1	TTTTACTTCG	TACTTGTGTT	TGCTTAAACA	AAGTGACTTG	TTTGGCTTAT
	humanEGR1
		4301				4350
30	mouseEGR1
	ratEGR1	AAACACATTG	AATGCGCTTT	ACTGCCCATG	GGATATGTGG	TGTGTATCCT
	humanEGR1
		4351			4388	
35	mouseEGR1
	ratEGR1	TCAGAAAAAT	TAAAAGGAAA	ATAAAGAAAC	TAAGTGGT	
	humanEGR1

Culture conditions and DNAzyme transfection. Primary rat aortic SMCs were obtained from Cell Applications, Inc., and grown in Waymouth's medium, pH 7.4, containing 10% fetal bovine serum (FBS), 50 μ g/ml streptomycin and 50 IU/ml penicillin at 37 °C in a humidified atmosphere of 5% CO₂. SMCs were used in experiments between passages 3-7. Pup rat SMCs (WKY12-22 (as described in Lemire et al, 1994, the entire contents of which are incorporated herein by reference)) were grown under similar conditions. Subconfluent (60-70%) SMCs were incubated in serum-free medium (SFM) for 6 h prior to DNAzyme (or antisense ODN, where indicated) transfection (0.1 μ M) using Superfect in accordance with manufacturer's instructions (Qiagen). After 18 h, the cells were washed with phosphate-buffered saline (PBS). pH 7.4 prior to transfection a second time in 5% FBS.

Northern blot analysis. Total RNA was isolated using the TRIzol reagent (Life Technologies) and 25 μ g was resolved by electrophoresis prior to transfer to Hybond-N+ membranes (NEN-DuPont). Prehybridization, hybridization with α^{32} P-dCTP-labeled Egr-1 or β -Actin cDNA, and washing

5 was performed essentially as previously described (Khachigian et al, 1995).

Western blot analysis. Growth-quiescent SMCs in 100 mm plates (Nunc-InterMed) were transfected with ED5 or ED5SCR as above, and incubated with 5% FBS for 1 h. The cells were washed in cold PBS, pH 7.4, and extracted in 150 mM NaCl, 50 mM Tris-HCl, pH 7.5, 1% sodium

10 deoxycholate, 0.1% SDS, 1% Triton X-100, 5 mM EDTA, 1% trasylol, 10 μ g/ml leupeptin, 1% aprotinin and 2 mM PMSF. Twenty four μ g protein samples were loaded onto 10% denaturing SDS-polyacrylamide gels and electroblotted onto PVDF nylon membranes (NEN-DuPont). Membranes were air dried prior to blocking with non-fat skim milk powder in PBS containing

15 0.05% (w:v) Tween 20. Membranes were incubated with rabbit antibodies to Egr-1 or Sp1 (Santa Cruz Biotechnology, Inc.) (1:1000) then with HRP-linked mouse anti-rabbit Ig secondary antiserum (1:2000). Where mouse monoclonal c-Fos (Santa Cruz Biotechnology, Inc.) was used, detection was achieved with HRP-linked rabbit anti-mouse Ig. Proteins were visualized by

20 chemiluminescent detection (NEN-DuPont).

Assays of cell proliferation. Growth-quiescent SMCs in 96-well titer plates (Nunc-InterMed) were transfected with ED5 or ED5SCR as above, then exposed to 5% FBS at 37 °C for 72 h. The cells were rinsed with PBS, pH 7.4, trypsinized and the suspension was quantitated using an automated Coulter

25 counter.

Assessment of DNase stability. DNases were 5'-end labeled with γ^{32} P-dATP and separated from free label by centrifugation. Radiolabeled DNases were incubated in 5% FBS or serum-free medium at 37 °C for the times indicated. Aliquots of the reaction were quenched by transfer to tubes

30 containing formamide loading buffer (Sambrook et al, 1989). Samples were applied to 12% denaturing polyacrylamide gels and autoradiographed overnight at -80 °C.

SMC wounding assay. Confluent growth-quiescent SMCs in chamber slides (Nunc-InterMed) were exposed to ED5 or ED5SCR for 18 h prior to a

35 single scrape with a sterile toothpick. Cells were treated with mitomycin C (Sigma) (20 μ M) for 2 h prior to injury (Pitsch et al, 1996; Horodyski &

Powell, 1996). Seventy-two h after injury, the cells were washed with PBS, pH 7.4, fixed with formaldehyde then stained with hematoxylin-eosin.

Rat arterial ligation model and analysis. Adult male Sprague Dawley rats weighing 300-350 g were anaesthetised using ketamine (60 mg/kg, i.p.) and xylazine (8 mg/kg, i.p.). The right common carotid artery was exposed up to the carotid bifurcation via a midline neck incision. Size 6/0 non-absorbable suture was tied around the common carotid proximal to the bifurcation, ensuring cessation of blood flow distally. A 200 μ l solution at 4°C containing 500 μ g of DNAzyme (in DEPC-treated H₂O), 30 μ l of transfecting agent and Pluronic gel P127 (BASF) was applied around the vessel in each group of 5 rats, extending proximally from the ligation for 12-15 mm. These agents did not inhibit the solidification of the gel at 37 °C. After 3 days, vehicle with or without 500 μ g of DNAzyme was administered a second time. Animals were sacrificed 18 days after ligation by lethal injection of phenobarbitone, and perfusion fixed using 10% (v:v) formaldehyde perfused at 120 mm Hg. Both carotids were then dissected free and placed in 10% formaldehyde, cut in 2 mm lengths and embedded in 3% (w:v) agarose prior to fixation in paraffin. Five μ m sections were prepared at 250 μ m intervals along the vessel from the point of ligation and stained with hematoxylin and eosin. The neointimal and medial areas of 5 consecutive sections per rat were determined digitally using a customized software package (Magellan) (Halasz & Martin, 1984) and expressed as a mean ratio per group of 5 rats.

Results and Discussion

The 7x7 nt arms flanking the 15 nt DNAzyme catalytic domain in the original DNAzyme design 7 were extended by 2 nts per arm for improved specificity (L.-Q. Sun, data not shown) (Figure 1). The 3' terminus of the molecule was capped with an inverted 3'-3'-linked thymidine (T) to confer resistance to 3'→5' exonuclease digestion. The sequence in both arms of ED5 was scrambled (SCR) without altering the catalytic domain to produce DNAzyme ED5SCR (Figure 1).

A synthetic RNA substrate comprised of 23 nts, matching nts 805 to 827 of NGFI-A mRNA (Figure 1) was used to determine whether ED5 had the capacity to cleave target RNA. ED5 cleaved the ³²P-5'-end labeled 23-mer within 10 min. The 12-mer product corresponds to the length between the

A(816)-U(817) junction and the 5' end of the substrate (Figure 1). In contrast, ED5SCR had no demonstrable effect on this synthetic substrate. Specific ED5 catalysis was further demonstrated by the inability of the human equivalent of this DNAzyme (hED5) to cleave the rat substrate over a wide range of stoichiometric ratios. Similar results were obtained using ED5SCR (data not shown): hED5 differs from the rat ED5 sequence by 3 of 18 nts in its hybridizing arms (Table 2). The catalytic effect of ED5 on a ³²P-labeled 206 nt fragment of native NGFI-A mRNA prepared by *in vitro* transcription was then determined. The cleavage reaction produced two radiolabeled species of 163 and 43 nt length consistent with DNAzyme cleavage at the A(816)-U(817) junction. In other experiments, ED5 also cleaved a ³²P-labeled NGFI-A transcript of 1960 nt length in a specific and time-dependent manner (data not shown).

Table 2. DNzyme target sites in mRNA.

Similarity between the 18 nt arms of ED5 or hED5 and the mRNA of rat NGFI-A or human EGR-1 (among other transcription factors) is expressed as a percentage. The target sequence of ED5 in NGFI-A mRNA is 5'-807-A CGU CCG GGA UGG CAG CGG-825-3' (SEQ ID NO: 13) (rat NGFI-A sequence), and that of hED5 in EGR-1 is 5'-262-U CGU CCA GGA UGG CCG CGG-280-3' (SEQ ID NO: 14) (Human EGR-1 sequence). Nucleotides in bold indicate mismatches between rat and human sequences. Data obtained by a gap best fit search in ANGIS using sequences derived from Genbank and EMBL. Rat sequences for Sp1 and c-Fos have not been reported.

	Gene	Accession number	Best homology over 18 nts (%)	
			ED5	hED5
15				
	Rat NGFI-A	M18416	100	84.2
20	Human EGR-1	X52541	84.2	100
	Murine Sp1	AF022363	66.7	66.7
	Human c-Fos	K00650	66.7	66.7
	Murine c-Fos	X06769	61.1	66.7
	Human Sp1	AF044026	38.9	28.9
25				

To determine the effect of the DNAzymes on endogenous levels of NGFI-A mRNA, growth-quiescent SMCs were exposed to ED5 prior to stimulation with serum. Northern blot and densitometric analysis revealed that ED5 (0.1 μ M) inhibited serum-inducible steady-state NGFI-A mRNA levels by 55% (Figure 2a), whereas ED5SCR had no effect (Figure 2a). The capacity of ED5 to inhibit NGFI-A synthesis at the level of protein was assessed by Western blot analysis. Serum-induction of NGFI-A protein was suppressed by ED5. In contrast, neither ED5SCR nor EDC, a DNAzyme bearing an identical catalytic domain as ED5 and ED5SCR but flanked by nonsense arms had any influence on the induction of NGFI-A (data not

shown). ED5 failed to affect levels of the constitutively expressed, structurally -related zinc-finger protein, Sp1. It was also unable to block serum-induction of the immediate-early gene product, c-Fos whose induction, like NGFI-A, is dependent upon serum response elements in its promoter and phosphorylation mediated by extracellular-signal regulated kinase (Treisman, 1990, 1994 and 1995; Gashler & Sukhatme, 1995). These findings, taken together, demonstrate the capacity of ED5 to inhibit production of NGFI-A mRNA and protein in a gene-specific and sequence-specific manner, consistent with the lack of significant homology between its target site in NGFI-A mRNA and other mRNA (Table 2).

The effect of ED5 on SMC replication was then determined. Growth-quiescent SMCs were incubated with DNase prior to exposure to serum and the assessment of cell numbers after 3 days. ED5 (0.1 μ M) inhibited SMC proliferation stimulated by serum by 70% (Figure 3a). In contrast, ED5SCR failed to influence SMC growth (Figure 3a). AS2, an antisense NGFI-A ODN able to inhibit SMC growth at 1 μ M failed to inhibit proliferation at the lower concentration (Figure 3a). Additional experiments revealed that ED5 also blocked serum-inducible 3 H-thymidine incorporation into DNA (data not shown). ED5 inhibition was not a consequence of cell death since no change in morphology was observed, and the proportion of cells incorporating Trypan Blue in the presence of serum was not influenced by either DNase (Figure 3b).

Cultured SMCs derived from the aortae of 2 week-old rats (WKY12-22) are morphologically and phenotypically similar to SMCs derived from the neointima of balloon-injured rat arteries (Seifert et al, 1984; Majesky et al, 1992). The epitheloid appearance of both WKY12-22 cells and neointimal cells contrasts with the elongated, bipolar nature of SMCs derived from normal quiescent media (Majesky et al, 1988). WKY12-22 cells grow more rapidly than medial SMCs and overexpress a large number of growth-regulatory molecules (Lemire et al, 1994), such as NGFI-A (Rafty & Khachigian, 1998), consistent with a "synthetic" phenotype (Majesky et al, 1992; Campbell & Campbell, 1985). ED5 attenuated serum-inducible WKY12-22 proliferation by approximately 75% (Figure 3c). ED5SCR had no inhibitory effect; surprisingly, it appeared to stimulate growth (Figure 3c). Trypan Blue exclusion revealed that DNase inhibition was not a consequence of cytotoxicity (data not shown).

To ensure that differences in the biological effects of ED5 and ED5SCR were not the consequence of dissimilar intracellular localization, both DNazymes were 5'-end labeled with fluorescein isothiocyanate (FITC) and incubated with SMCs. Fluorescence microscopy revealed that both FITC-ED5 and FITC-ED5SCR localized mainly within the nuclei. Punctate fluorescence in this cellular compartment was independent of DNzyme sequence. Fluorescence was also observed in the cytoplasm, albeit with less intensity. Cultures not exposed to DNzyme showed no evidence of autofluorescence.

Both molecules were 5'-end labeled with $\gamma^{32}\text{P}$ -dATP and incubated in culture medium to ascertain whether cellular responsiveness to ED5 and ED5SCR was a consequence of differences in DNAzyme stability. Both ^{32}P -ED5 and ^{32}P -ED5SCR remained intact even after 48 h. In contrast to ^{32}P -ED5 bearing the 3' inverted T, degradation of ^{32}P -ED5 bearing its 3' T in the correct orientation was observed as early as 1 h. Exposure to serum-free medium did not result in degradation of the molecule even after 48 h. These findings indicate that inverse orientation of the 3' base in the DNAzyme protects the molecule from nucleolytic cleavage by components in serum.

Physical trauma imparted to SMCs in culture results in outward migration from the wound edge and proliferation in the denuded zone. We determined whether ED5 could modulate this response to injury by exposing growth-quiescent SMCs to either DNazyme and Mitomycin C, an inhibitor of proliferation (Pitsch et al, 1996; Horodyski & Powell, 1996) prior to scraping. Cultures in which DNazyme was absent repopulated the entire denuded zone within 3 days. ED5 inhibited this reparative response to injury and prevented additional growth in this area even after 6 days (data not shown). That ED5SCR had no effect in this system further demonstrates sequence-specific inhibition by ED5.

The effect of ED5 on neointima formation was investigated in a rat model. Complete ligation of the right common carotid artery proximal to the bifurcation results in migration of SMCs from the media to the intima where proliferation eventually leads to the formation of a neointima (Kumar & Lindner, 1997; Bhawan et al, 1977; Buck, 1961). Intimal thickening 18 days after ligation was inhibited 50% by ED5 (Figure 4). In contrast, neither its scrambled counterpart (Figure 4) nor the vehicle control (Figure 4) had any effect on neointima formation. These findings demonstrate the capacity of

ED5 to suppress SMC accumulation in the vascular lumen in a specific manner, and argue against inhibition as a mere consequence of a "mass effect" (Kitze et al, 1998; Tharlow et al, 1996).

Further experiments revealed the capacity of hED5 to cleave (human) EGR-1 RNA. hED5 cleaved its substrate in a dose-dependent manner over a wide range of stoichiometric ratios. hED5 also cleaved in a time-dependent manner, whereas hED5SCR, its scrambled counterpart, had no such catalytic property (data not shown).

The specific, growth-inhibitory properties of ED5 reported herein suggest that DNazymes may be useful as therapeutic tools in the treatment of vascular disorders involving inappropriate SMC growth.

Example 2

Cleavage of human EGR-1 RNA by panel of candidate DNazymes

To evaluate which specific DNazymes targeting human EGR-1 (other than hED5) efficiently cleave EGR-1 RNA, we prepared *in vitro* transcribed 35S-labeled EGR-1 RNA and incubated this substrate with candidate DNazymes for various times. The EGR-1 plasmid template (hs164) was prepared by subcloning bps 168-332 of human EGR-1 into pGEM-T-easy. A 388 nt 35S-labeled substrate was prepared by *in vitro* transcription using SP6 polymerase. Time-dependent cleavage of the substrate was tested using the following DNazymes:

- DzA: 5'-CAGGGGACAGGCTAGCTACAACGACGTTGCGGG-X-3' (SEQ ID NO: 15) ;
 - DzB: 5'-TGCAGGGGAGGCTAGCTACAACGAACCGTTGCG-X-3' (SEQ ID NO: 16) ;
 - DzC: 5'-CATCCTGGAGGCTAGCTACAACGAGAGCAGGCT-X-3' (SEQ ID NO: 17) ;
 - DzE: 5'-TCAGCTGCAGGCTAGCTACAACGACTCGGCCTT-X-3' (SEQ ID NO: 18) ; and
 - DzF: 5'-GCGGGGACAGGCTAGCTACAACGACAGCTGCAT-X-3' (SEQ ID NO: 19)
- where X denotes a 3'-3-linked T.

The porcine and human EGR-1 sequences are remarkably well conserved (91%). Porcine retinal SMCs were used to determine whether DzA could block the growth of porcine SMCs. Our studies indicate that DzA (0.5 μ M) could inhibit the proliferation of these cells (Figure 6). In contrast, DzE had no effect (Figure 6).

Example 6

Delivery of DNAzyme into the porcine coronary artery wall

5 Porcine angioplasty and stenting are accepted models of human in-
 stent restenosis (Karas et al. 1992). The porcine coronary anatomy,
 dimensions and histological response to stenting are similar to the human
 (Muller et al. 1992). The Transport Catheter has previously been used to
 deliver antisense DNA targeting c-myc in humans (Serrys et al.
 10 1998) and the pig (Gunn & Cumberland, 1996) via the intraluminal route.
 Using this catheter, FITC-labeled DNAzyme was applied to the inner wall of
 a porcine coronary artery, *ex vivo*, from a newly explanted pig heart.
 DNAzyme (1000 µg) was delivered via the catheter in 2ml MilliQ H2O
 containing 300µl FuGENE6 and 1mM MgCl₂. The FITC-labeled DNAzyme
 15 localised into the intimal cells of the vessel wall. These studies demonstrate
 that DNAzyme can be delivered to cells within the artery wall via an
 intraluminal catheter.

Throughout this specification the word "comprise", or variations such
 as "comprises" or "comprising", will be understood to imply the inclusion of a
 20 stated element, integer or step, or group of elements, integers or steps, but
 not the exclusion of any other element, integer or step, or group of elements,
 integers or steps.

It will be appreciated by persons skilled in the art that numerous
 variations and/or modifications may be made to the invention as shown in
 25 the specific embodiments without departing from the spirit or scope of the
 invention as broadly described. The present embodiments are, therefore, to
 be considered in all respects as illustrative and not restrictive. In addition,
 various documents are cited throughout this application. The disclosures of
 these documents are hereby incorporated by reference into this application to
 30 describe more fully the state of the art to which this invention pertains.

References

- Autieri, M.V. et al (1995) Antisense oligonucleotides to the P65 subunit of NF- κ B inhibit human vascular smooth muscle cell adherence and proliferation and prevent neointima formation in rat carotid arteries. *Biochem. and Biophys. Res. Commun.* 213:827-836.
- Bennett, M.D. and Schwartz, S.M. (1995) Antisense Therapy for Angioplasty Restenosis. *Circulation* 92:1981-1993.
- Bhawan, J., Joris, I., DeGerolami, U. & Majno, G. (1977) Effect of occlusion on large vessels. *Am. J. Pathol.* 88, 355-380.
- Breaker, R.R. and Joyce, G. (1994) *Chemistry and Biology* 1:223-229.
- Breaker, R.R. and Joyce, G. (1995) *Chemistry and Biology* 2:655-660.
- Brogi, E., et al. (1993) Distinct patterns of expression of fibroblast growth factors and their receptors in human atheroma and nonatherosclerotic arteries. *J. Clin. Invest.* 92, 2408-2418.
- Buck, R.C. (1961) Intimal thickening after ligation of arteries. *Circ. Res.* 9, 418-426.
- Campbell, G.R. & Campbell, J.H. (1985) Smooth muscle phenotypic changes in arterial wall homeostasis: implications for the pathogenesis of atherosclerosis. *Exp. Mol. Pathol.* 42, 139-162.
- Carmi, N., et al. (1996) *Chemistry and Biology* 3:1039-1046.
- Delbridge, G.J. & Khachigian, L.M. (1997) FGF-1-induced PDGF A-chain gene expression in vascular endothelial cells involves transcriptional activation by Egr-1. *Circ. Res.* 81, 282-288.

- Evanko, S.P., Raines, E.W., Ross, R., Gold, L.I. & Wight, T.N. (1998) Proteoglycan distribution in lesions of atherosclerosis depends on lesion severity, structural characteristics, and the proximity of platelet-derived growth factor and transforming growth factor-beta. Am. J. Pathol. 152, 533-546.
5
- Frimerman, A. et al. (1999) Chimeric DNA-RNA Hammerhead Ribozyme to Proliferating Cell Nuclear Antigen Reduces Stent-Induced Stenosis in a Porcine Coronary Model. Circulation 99: 697-703.
10
- Gashler, A. & Sukhatme, V. (1995) Early growth response protein 1 (Egr-1): prototype of a zinc-finger family of transcription factors. Prog. Nucl. Acid Res. 50, 191-224.
- 15 Gunn J. and Cumberland D.C. (1996) Dual balloon catheter: seminars in Interventional Cardiology 1:31-33.
- Halasz, P. & Martin, P. (1984) A microcomputer-based system for semi-automatic analysis of histological sections. Proc. Royal Microscop. Soc. 19, 312.
20
- Haseloff, J. & Gerlach, W.A. (1988) Simple RNA enzymes with new and highly specific endoribonuclease activities. Nature 334, 585-591.
- 25 Holmes, D.R., et al. (1984) Restenosis after percutaneous transluminal coronary angioplasty (PTCA): a report from the PTCA Registry of the National Heart, Lung, and Blood Institute. Am. J. Cardiol. 53, 77C-81C.
- Horodyski, J. & Powell, R.J. (1996) Effect of aprotinin on smooth muscle cell proliferation, migration, and extracellular matrix synthesis. J. Surg. Res. 66, 115-118.
30
- Horodyski, J. & Powell, R.J. (1996) Effect of aprotinin on smooth muscle cell proliferation, migration, and extracellular matrix synthesis. J. Surg. Res. 66, 115-118.
35

- Hughes, S.E., Crossman, D. & Hall, P.A. (1993) Expression of basic and acidic fibroblast growth factors and their receptor in normal and atherosclerotic human arteries. *Cardiovasc. Res.* 27, 1214-1219.
- 5 Jackson, C.L. & Schwartz, S.M. (1992) Pharmacology of smooth muscle replication. *Hypertension* 20, 713-736.
- Karas et al. (1992) Coronary intimal proliferation after balloon injury and stenting in the sine: an animal model of restenosis. *J. Am. Coll. Cardiol.* 20, 467-474.
- 10 Karas et al. (1992) Coronary intimal proliferation after balloon injury and stenting in the sine: an animal model of restenosis. *J. Am. Coll. Cardiol.* 20, 467-474.
- Kashani-Sabet, M., et al. (1992) Antisense Research and Development 2:3-15.
- 15 Khachigian, L.M. & Collins, T. (1997) Inducible expression of Egr-1-dependent genes: a paradigm of transcriptional activation in vascular endothelium. *Circ. Res.* 81, 457-461.
- Khachigian, L.M., et al. (1997) Egr-1 is activated in endothelial cells exposed to fluid shear stress and interacts with a novel shear-stress response element in the PDGF A-chain promoter. *Arterioscl. Thromb. Vasc. Biol.* 17, 2280-2286.
- 20 Khachigian, L.M., et al. (1997) Egr-1 is activated in endothelial cells exposed to fluid shear stress and interacts with a novel shear-stress response element in the PDGF A-chain promoter. *Arterioscl. Thromb. Vasc. Biol.* 17, 2280-2286.
- Khachigian, L.M., Lindner, V., Williams, A.J. & Collins, T. (1996) Egr-1-induced endothelial gene expression: a common theme in vascular injury. *Science* 271, 1427-1431.
- 25 Khachigian, L.M., Lindner, V., Williams, A.J. & Collins, T. (1996) Egr-1-induced endothelial gene expression: a common theme in vascular injury. *Science* 271, 1427-1431.
- Khachigian, L.M., Williams, A.J. & Collins, T. (1995) Interplay of Sp1 and Egr-1 in the proximal PDGF-A promoter in cultured vascular endothelial cells. *J. Biol. Chem.* 270, 27679-27686.
- 30 Kim, S., et al. (1995) Angiotensin II type 1 receptor blockade inhibits the expression of immediate-early genes and fibronectin in rat injured artery. *Circulation* 92, 88-95.

- Kitze, B., et al. (1998) Human CD4+ T lymphocytes recognize a highly conserved epitope of human T lymphotropic virus type 1 (HTLV-1) env gp21 restricted by HLA DRB1*0101. Clin. Exp. Immunol. 111, 278-285.
- 5 Koizumi, M., et al. (1989) Nucleic Acids Research 17:7059-7069.
- Kumar, A. & Lindner, V. (1997) Remodeling with neointima formation in the mouse carotid artery after cessation of blood flow. Arterioscl. Thromb. Vasc. Biol. 17, 2238-2244.
- 10 Lemire, J.M., Covin, C.W., White, S., Giachelli, C.M. & Schwartz, S.M. (1994) Characterization of cloned aortic smooth muscle cells from young rats. Am. J. Pathol. 144, 1068-1081.
- 15 Libby, P., Schwartz, S.M., Brogi, E., Tanaka, H. & Clinton, S. (1995) A cascade model for restenosis. Circ. Res. 86 (Suppl. III), 47-52.
- Majesky, M.W., Benditt, E.P. & Schwartz, S.M. (1988) Expression and developmental control of platelet-derived growth factor A-chain and B-chain/Sis genes in rat aortic smooth muscle cells. Proc. Natl. Acad. Sci. USA 20 85, 1524-1528.
- Majesky, M.W., Giachelli, C.M., Reidy, M.A. & Schwartz, S.M. (1992) Rat carotid neointimal smooth muscle cells reexpress a developmentally 25 regulated mRNA phenotype during repair of arterial injury. Circ. Res. 71, 759-768.
- Milbrandt, J. (1987) A nerve growth factor-induced gene encodes a possible transcriptional regulatory factor. Science 238, 797-799
- 30 Morishita, R. et al (1993) Single intraluminal delivery of antisense cdc2 kinase and proliferating-cell nuclear antigen oligonucleotides result in chronic inhibition of neointimal hyperplasia. Proc. Natl. Acad. Sci. USA 90:8474-8478.

Muller DW et al. (1992) Experimental models of coronary artery restenosis. *J. Am. Coll. Cardiol.* 19, 418-432.

Murry, C.E., Bartosek, T., Giachelli, C.M., Alpers, C.E. & Schwartz, S.M. (1996) Platelet-derived growth factor-A mRNA expression in fetal, normal adult, and atherosclerotic human aortas. *Circulation* 93, 1095-1106

Otsuka, E. and Koizumi, M., Japanese Patent No. 4,235,919.

Pan, T. and Uhlenbeck, O.C. (1996) *Biochemistry* 31:3887-3895.

Pitsch, R.J., et al. (1996) Inhibition of smooth muscle cell proliferation and migration in vitro by antisense oligonucleotide to c-myb. *J. Vasc. Surg.* 23, 783-791.

Pitsch, R.J., et al. (1996) Inhibition of smooth muscle cell proliferation and migration in vitro by antisense oligonucleotide to c-myb. *J. Vasc. Surg.* 23, 783-791.

Raillard, S.A. and Joyce, G.F. (1996) *Biochemistry* 35:11693-11701.

Rafty, L.A. & Khachigian, L.M. (1998) Zinc finger transcription factors mediate high constitutive PDGF-B expression in smooth muscle cells derived from aortae of newborn rats. *J. Biol. Chem.* 273, 5758-5764.

Rekhter, M. & Gordon, D. (1994) Does platelet-derived growth factor-A chain stimulate proliferation of arterial mesenchymal cells in human atherosclerotic plaques? *Circ. Res.* 75, 410-417.

Sambrook, J., Fritsch, E.F. & Maniatis, T. (1989) *Molecular cloning: a laboratory manual*, Cold Spring Harbor Laboratory Press, Plainview, N.Y.

Santoro, S.W. & Joyce, G.F. (1997) A general purpose RNA-cleaving DNA enzyme. *Proc. Natl. Acad. Sci. USA* 94, 4262-4266.

Seifert, R.A., Schwartz, S.M. & Bowen-Pope, D.F. (1984) Developmentally regulated production of platelet-derived growth factor-like molecules. *Nature* 311, 669-671.

5 Serrys PW et al. (1998) Antisense oligonucleotide against *c-myc* administered with the Transport Catheter (36-48 holes, 250 μ m diameter) for the prevention of in-stent restenosis: results of randomised ITALICS trial. *Circulation* :11909.

10 Silverman, E.S., Khachigian, L.M., Lindner, V., Williams, A.J. & Collins, T. (1997) Inducible PDGF A-chain transcription in vascular smooth muscle cells is mediated by Egr-1 displacement of Sp1 and Sp3. *Am. J. Physiol.* 42, H1415-H1426.

15 Stary, H.C., et al. (1995) A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Atherosclerosis, American Heart Association. *Arterioscler. Thromb. Vasc. Biol.* 15, 1512-1531.

20 Sumpio, B.E., et al. (1998) Regulation of PDGF-B by cyclic strain: lack of involvement of the shear-stress responsive element. *Arterioscler. Thromb. Vasc. Biol.* 18, 349-355.

Sun, L.Q., et al. (1997) *Mol. Biotechnology* 7:241-251.

25 Simons, M. et al (1992) Antisense *c-myb* oligonucleotides inhibit intimal arterial smooth muscle cell accumulation *in vivo*. *Nature* 359:67-70.

Symonds, R.H. (1992) *Ann. Rev. Biochem.* 61:641-671.

30 Tanizawa, S., Ueda, M., van der Loos, C.M., van der Wal, A.C. & Becker, A.E. (1996) Expression of platelet-derived growth factor B-chain and beta-receptor expression in human coronary arteries after percutaneous transluminal coronary angioplasty: an immunohistochemical study. *Heart* 75, 549-556

- Tharlow, R.J., Hill, D.R. & Woodruff, G.N. (1996) Comparison of the autoradiographic binding distribution of [3H]-gabapentin with excitatory amino acid receptor and amino acid uptake site distributions in rat brain. *Brit. J. Pharmacol.* 118, 457-465.
- 5 Treisman, R. (1995) Journey to the surface of the cell: Fos regulation and the SRE. *EMBO J.* 14, 4905-4913.
- Treisman, R. (1994) Ternary complex factor: growth factor regulated transcriptional activators. *Curr. Opin. Genet. Develop.* 4, 96-101.
- 10 Treisman, R. (1990) The SRE: a growth factor responsive transcriptional regulator. *Sem. Cancer Biol.* 1, 47-58.
- 15 Tsang, J. and Joyce, G.F. (1994) *Biochemistry* 33:5966-5973.
- Ueda, M., et al. (1996) In situ detection of platelet-derived growth factor-A and -B chain mRNA in human coronary arteries after percutaneous transluminal coronary angioplasty. *Am. J. Pathol.* 149, 831-843.
- 20 Wilcox, J.N., Smith, K.M., Schwartz, S.M. & Gordon, D. (1989) Localization of tissue factor in the normal vessel wall and in the atherosclerotic plaque. *Proc. Natl. Acad. Sci. USA* 86, 2839-2843.
- 25 Wilcox, J.N., Smith, K.M., Williams, L.T., Schwartz, S.M. & Gordon, D. (1988) Platelet-derived growth factor mRNA detection in human atherosclerotic plaques by in situ hybridization. *J. Clin. Invest.* 82, 1134-1143.

1. A DNzyme which specifically cleaves EGR-1 mRNA, the DNzyme comprising

- (i) a catalytic domain which cleaves mRNA at a purine:pyrimidine cleavage site;
- (ii) a first binding domain contiguous with the 5' end of the catalytic domain; and

- 10 (iii) a second binding domain contiguous with the 3' end of the catalytic domain,

wherein the binding domains are sufficiently complementary to the two regions immediately flanking a purine:pyrimidine cleavage site within the region of EGR-1 mRNA corresponding to nucleotides 168 to 332 as shown in SEQ ID NO:1, such that the DNAzyme cleaves the EGR-1 mRNA.

2. A DNzyme as claimed in claim 1 wherein each binding domain is nine or more nucleotides in length.

- 20 3. A DNAzyme as claimed in claim 1 or claim 2 in which the catalytic domain has the nucleotide sequence GGCTAGCTACAACGA (SEQ ID NO: 2).

4. A DNazyme as claimed in any one of claims 1 to 3 in which the cleavage site is selected from the group consisting of

- 25 (i) the GU site corresponding to nucleotides 198-199;
(ii) the GU site corresponding to nucleotides 200-201;
(iii) the GU site corresponding to nucleotides 264-265;
(iv) the AU site corresponding to nucleotides 271-272;
(v) the AU site corresponding to nucleotides 301-302;
30 (vi) the GU site corresponding to nucleotides 303-304; and
(vii) the AU site corresponding to nucleotides 316-317.

5. A DNAzyme as claimed in claim 4 in which the cleavage site is the AU site corresponding to nucleotides 271-272.

6. A DNzyme as claimed in claim 1 which has a sequence selected from the group consisting of:

- 5 (i) 5'-caggggacaGGCTAGCTACAACGAcgttgcg (SEQ ID NO: 3);
- (ii) 5'-tgcaggggaGGCTAGCTACAACGAaccgttgcg (SEQ ID NO: 4);
- (iii) 5'-catcctggaGGCTAGCTACAACGAgagcaggct (SEQ ID NO: 5);
- (iv) 5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6);
- (v) 5'-ccgctgccaGGCTAGCTACAACGAcccggacgt (SEQ ID NO: 7);
- 10 (vi) 5'-gcggggacaGGCTAGCTACAACGAcagctgcat (SEQ ID NO: 8);
- (vii) 5'-cagcggggaGGCTAGCTACAACGAatcagctgc (SEQ ID NO: 9); and
- (viii) 5'-ggtcagagaGGCTAGCTACAACGActgcagcg (SEQ ID NO: 10).

15 7. A DNzyme as claimed in claim 6 which has the sequence:
5'-ccgcggccaGGCTAGCTACAACGAcctggacga (SEQ ID NO: 6).

8. A DNzyme as claimed in any one of claims 1 to 7, wherein the 3'-end nucleotide residue is inverted in the binding domain contiguous with the 3' end of the catalytic domain.

9. A pharmaceutical composition comprising a DNzyme according to any one of claims 1 to 8 and a pharmaceutically acceptable carrier.

25 10. A method of inhibiting EGR-1 activity in cells which comprises exposing the cells to a DNzyme according to any one of claims 1 to 8.

11. A method of inhibiting proliferation or migration of cells in a subject which comprises administering to the subject a prophylactically effective dose of the pharmaceutical composition according to claim 9.

12. A method of treating a condition associated with cell proliferation or migration in a subject which comprises administering to the subject a therapeutically effective dose of the pharmaceutical composition according to claim 9.

19. A method for inhibiting the onset of restenosis in a subject undergoing angioplasty, which comprises topically administering a stent according to claim 15 to the subject at around the time of the angioplasty.

WO 00/42173

PCT/AU00/00011

1/8





Figure 1

2/8




Figure 2




Figure 3A




Figure 3B




Figure 3C




Figure 4




Figure 5





Figure 6



DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name. I believe that I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled "CATALYTIC MOLECULES," the specification of which was filed as International Application No. PCT/AU00/00011 on January 11, 2000 and was transmitted to the U.S. Receiving Office on July 11, 2001 together with a Preliminary Amendment dated July 11, 2001, and which was assigned Application Serial No. 09/889,075. I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

			Priority Claimed
<u>PP8103</u>	<u>Australia</u>	<u>11/January/1999</u>	<input checked="" type="checkbox"/> <input type="checkbox"/>
(Application Serial Number)	(Country)	(Day/Month/Year Filed)	Yes No

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below:

<u>NONE</u>	
(Application Serial Number)	(Day/Month/Year Filed)

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) or PCT international application(s) designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior application(s) in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

<u>NONE</u>		
(Application Serial Number)	(Day/Month/Year Filed)	(Status-Patented, Pending or Abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: I hereby appoint as my attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

John B. Lungmus (18,566)
Allen H. Gerstein (22,218)
Nate F. Scarpelli (22,320)
Michael F. Borun (25,447)
Trevor B. Joike (25,542)
Carl E. Moore, Jr. (26,487)


Richard H. Anderson (26,526)
Patrick D. Ertel (26,877)
Richard B. Hoffman (26,910)
James P. Zeller (28,491)
Kevin D. Hogg (31,839)
Jeffrey S. Sharp (31,879)


Martin J. Hirsch (32,237)
James J. Napoli (32,361)
Richard M. La Barge (32,254)
Douglass C. Hochstetler (33,710)
Robert M. Gerstein (34,824)
Anthony G. Sitko (36,278)


James A. Flight (37,622)
Roger A. Heppermann (37,641)
David A. Gass (38,153)
Gregory C. Mayer (38,238)
Michael R. Weiner (38,359)
William K. Merkel (40,725)

Send correspondence to: David A. Gass

FIRM NAME	PHONE NO.	STREET	CITY & STATE	ZIP CODE
Marshall, Gerstein & Borun	312-474-6300	6300 Sears Tower 233 South Wacker Drive	Chicago, Illinois	60606-6402

Full Name of First or Sole Inventor <u>David G. Atkins</u>	Citizenship AU
Residence Address - Street 45W 60th St., Apt 17A	Post Office Address - Street 45W 60th St., Apt 17A
City (Zip) New York 10023 NY	City (Zip) New York 10023
State or Country New York	State or Country New York
Date <input checked="" type="checkbox"/> 1 NOVEMBER 2001	Signature <input checked="" type="checkbox"/> 

Second Joint Inventor, if any <u>Andrew R. Baker</u>	Citizenship AU
Residence Address - Street 3 Adelong Place	Post Office Address - Street 3 Adelong Place
City (Zip) Wahroonga 2076 WLS	City (Zip) Wahroonga 2076
State or Country New South Wales	State or Country New South Wales
Date <input checked="" type="checkbox"/> 15 NOVEMBER 2001	Signature <input checked="" type="checkbox"/> 

Third Joint Inventor, if any <u>Levon Michael Khachigian</u>	Citizenship AU
Residence Address - Street 5 Ratcliffe Street	Post Office Address - Street 5 Ratcliffe Street
City (Zip) Ryde 2112	City (Zip) Ryde 2112
State or Country New South Wales WLS	State or Country New South Wales
Date <input checked="" type="checkbox"/> 9 November 2001	Signature <input checked="" type="checkbox"/> 

WO 00/42173

PCT/AU00/00011

1/8

SEQUENCE LISTING

<110> Johnson & Johnson Pty Ltd
Unisearch Limited

5 <120> Catalytic molecules

<150> PP 8103
<151> 1999-01-11

10 <160> 20

<170> PatentIn Ver. 2.1

15 <210> 1
<211> 3132
<212> DNA
<213> Homo sapiens

20 <400> 1
ccgcagaact tggggagccg ccgcgcgccat ccgcgcgcgc agccagcttc cgcgcgcgca 60
ggaccggccc ctgccccagc ctccgcagcc gcggcgcgtc caccgccgcc cgcgcgccagg 120
gcgagtcggg gtgcgcgcct gcacgcttct cagtgttccc cgcgcgccgc atgtaaccgc 180
gccaggcccc cgcaacggtg tcccctgcag ctccagcccc gggctgcacc ccccgcccc 240
25 gacaccagct ctccagcctg ctctgtccagg atggcgcgcg ccaaggccga gatgcagctg 300
atgtccccgc tgcagatctc tgacccgttc ggatcctttc ctactcgcgc caccatggac 360
aactacccta agctggagga gatgatgctg ctgagcaacg gggctcccca gttcctcggc 420
gccgcgcggg ccccagaggg cagcggcagc aacagcagca gcagcagcag cggggggcgg 480
ggaggcggcg ggggcggcag caacagcagc agcagcagca gcaccttcaa cctcaggcgc 540
30 gacacgggcg agcagcccta cgagcacctg accgcagagt cttttcctga catctctctg 600
aacaacgaga aggtgctggt ggagaccagt taccacagcc aaaccactcg actgcccccc 660
atcacctata ctggcgcgtt ttccctggag cctgcaccca acagtggcaa cacttctgtg 720
cccgagcccc tcttcagctt ggtcagtggt ctagtgagca tgaccaaccc accggcctcc 780
tcgtcctcag caccatctcc agcggcctcc tccgcctccg cctcccagag cccaccctg 840
35 agctgcgcag tgccatccaa cgacagcagt cccatttact cagcggcacc cacttccccc 900
acgccgaaca ctgacatttt ccttgagcca caaagccagg ccttcccggg ctccggcagg 960
acagcgctcc agtaccgcgc tctgcctac cctgcgcgca aggggtggtt ccagggtccc 1020
atgatccccg actacctgtt tccacagcag cagggggatc tgggcctggg caccacagac 1080
cagaagccct tccagggcct ggagagccgc acccagcagc ctctcgtaac cctctgtct 1140
40 actattaagg cctttgccac tcagtcgggc tcccaggacc tgaaggccct caataccagc 1200
taccagtccc agctcatcaa acccagccgc atgcgcaagt atcccaaccg gccagcaag 1260
acgccccccc acgaacgccc ttacgcttgc ccagtggagt cctgtgatcg ccgcttctcc 1320
cgctccgacg agctcaccgc ccacatccgc atccacacag gccagaagcc ctccagtgcc 1380
cgcatctgca tgcgcaactt cagcgcgcgc gaccacctca ccaccacat ccgcacccac 1440
45 acaggcgaaa agcccttcgc ctgcgacatc tgtggaagaa agtttgccag gacgatgaa 1500
cgcaagaggc ataccaagat ccacttgccg cagaaggaca agaaagcaga caaaagtgtt 1560
gtggcctctt cggccacctc ctctctctct tccctaccgt ccccggttgc tactcttac 1620
ccgtcccccg ttactacctc ttatccatcc ccggccacca cctcatacc atccccctgt 1680
50 cccacctcct tctcctctcc cggctcctcg acctacccat cccctgtgca cagtggcttc 1740
ccctccccgt cgggtggccac cagctactcc tctgttcccc ctgctttccc ggcccaggtc 1800
agcagcttcc ctctcctcagc tgtcaccac tccttcagcg cctccacagg gctttcggac 1860
atgacagcaa ccttttctcc caggacaatt gaaatttgc aaagggaag gggaaagaaa 1920
gggaaaaggg agaaaaagaa acacaagaga cttaaaggac aggaggagga gatggccata 1980
ggagaggagg gtctctctta ggtcagatgg aggttctcag agccaagtcc tccctctcta 2040
55 ctggagtggg aggtctattg gccacaatc ctttctgccc acttcccctt ccccaattac 2100
tattcccttt gacttcagct gcctgaaaca gccatgtcca agttcttcac ctctatccaa 2160
agaacttgat ttgcatggat tttggataaa tcatttcagt atcatctcca tcatatgcct 2220
gacccttgc tcccttcaat gctagaaaat cgagttggca aaatgggggt tgggcccctc 2280
agagccctgc cctgcacctc tgtacagtgt ctgtgccatg gatttcgttt ttcttgggt 2340
60 actcttgatg tgaagataat ttgcatatc tattgtatta tttggagtta ggtcctcact 2400

<220>
<223> Description of Artificial Sequence: DNazyme

WO 00/42173

PCT/AU00/00011

3/8

<400> 5
catcctggag gctagctaca acgagagcag gct 33

5 <210> 6
<211> 33
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Description of Artificial Sequence: DNAzyme

<400> 6
ccgcggccag gctagctaca acgacctgga cga 33

15

<210> 7
<211> 33
<212> DNA
20 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: DNAzyme

25 <400> 7
ccgctgccag gctagctaca acgacccgga cgt 33

<210> 8
30 <211> 33
<212> DNA
<213> Artificial Sequence

<220>
35 <223> Description of Artificial Sequence: DNAzyme

<400> 8
gcggggacag gctagctaca acgacagctg cat 33

40

<210> 9
<211> 33
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence: DNAzyme

<400> 9
50 cagcggggag gctagctaca acgaatcagc tgc 33

<210> 10
<211> 33
55 <212> DNA
<213> Artificial Sequence

<220>
60 <223> Description of Artificial Sequence: DNAzyme

WO 00/42173

PCT/AU00/00011

4/8

<400> 10
ggtcagagag gctagctaca acgactgcag cgg

33

5 <210> 11
<211> 3068
<212> DNA
<213> Mus musculus

10 <400> 11
ggggagccgc cgcgcgcgatt cgcgcgcgcgc gccagcttcc gccgcgcgcaa gatcggccccc 60
tgccccagcc tccgcgcgcag ccctgcgtcc accacggggc gcggctaccg ccagcctggg 120
ggccccaccta cactccccgc agtgtgcccc tgcacccgcg atgtaaccgc gccaaccccc 180
ggcgagtgtg ccctcagtag ctccggcccc gggtgcgcgc caccacccaa catcagttct 240
15 ccagctcgct ggtccgggat ggcagcggcc aaggccgaga tgcaattgat gtctccgctg 300
cagatctctg acccgcttcgg ctcccttccct cactcaccga ccatggacaa ctaccccaaa 360
ctggaggaga tgatgctgct gagcaacggg gctccccagt tectcggtgc tgcgggaacc 420
ccagagggca gcggcggtta tagcagcagc agcaccagca gcggggggcg tgggtggggg 480
ggcagcaaca gcggcagcag cgccttcaat cctcaagggg agccgagcga acaaccctat 540
20 gagcacctga ccacagagtc cttttctgac atcgctctga ataagagaa ggcgatgggtg 600
gagacgagtt atcccagcca aacgactcgg ttgcctccca tcacctatac tggccgcttc 660
tccctggagc ccgcacccaa cagtggcaac actttgtggc ctgaacccct tttcagccta 720
gtcagtggcc tcgtgagcat gaccaatcct ccgacctctt catcctcggc gccttctcca 780
gctgcttcat cgtcttcttc tgcctcccag agcccgcccc tgagctgtgc cgtgcgctcc 840
25 aacgacagca gtcccatcta ctccgctgcg cccacctttc ctactcccaa cactgacatt 900
tttccctgagc cccaaagcca ggcctttcct ggctcggcag gcacagcctt gcagtaccgc 960
cctcctgect accctgccac caaagggtgg ttccaggttc ccatgatccc tgactatctg 1020
tttccacaac aacagggaga cctgagcctg ggcacccag accagaagcc cttccagggt 1080
ctggagaacc gtacccagca gccttcgctc actccactat ccactattaa agccttcgct 1140
30 actcagtcgg gctcccagga cttaaaggct cttaatacca cctaccaatc ccagctcatc 1200
aaaccagcc gcctgctgca gtaccccaac cggcccagca agacaccccc ccatgaacgc 1260
ccatatgctt gccctgtcga gtccctgcag cgcgctttt ctgctcggga tgagcttacc 1320
cgccatatcc gcctccacac aggcagaaag cccctccagt gtcgaatctg catgctgaac 1380
ttcagtcgta gtgaccacct taccacccac atccgcaccc acacaggcga gaagcctttt 1440
35 gcctgtgaca ttgtggggag gaagtttgcc aggagtgatg aacgcaagag gcataccaaa 1500
atccatttaa gacagaagga caagaaagca gacaaaagtg tgggtggcctc cccggctgcc 1560
tcttactctt cttcttacc atccccagtg gctacctctt accatcccc tgcacccacc 1620
tcattcccat cccctgtgcc cacttcctac tccctcctg gctcctccac ctacccatct 1680
cctgcgcaca gtggttccc gtcgcgctca gtggccacca cctttgcctc cgttccacct 1740
40 gctttcccca cccaggctcag cagcttcccg tctgcggcg tcagcagctc cttcagcacc 1800
tcaactggtc tttcagacat gacagcgacc ttttctccca ggacaattga aatttgctaa 1860
agggaataaa agaaagcaaa gggagaggca ggaaagacat aaaagcacag gaggaagag 1920
atggccgcaa gaggggccac ctcttaggtc agatggaaga tctcagagcc aagtccttct 1980
actcagagtg agaaggaccg ttggccaaca gccctttcac ttaccatccc tgcctcccc 2040
45 gtccgtgttc ctttgacttc agctgcctga aacagccatg tccaagtctt tcacctctat 2100
ccaaaggact tgatttgcat ggtattggat aaatcatttc agtatcctct ccatcacatg 2160
cctggccctt gctcccttca gcgctagacc atcaagttgg cataaagaaa aaaaaatggg 2220
tttgggccc cagaaccctg cctgcatct ttgtacagca tctgtgccat ggattttgtt 2280
ttccttgggg tattcttgat gtgaagataa ttgtcatact ctattgtatt atttgagatt 2340
50 aaatcctcac tttgggggag gggggagcaa agccaagcaa accaatgatg atcctctatt 2400
ttgtgatgac tctgctgtga cattaggttt gaagcatttt ttttttcaag cagcagtcct 2460
aggtattaac tggagcatgt gtcagagtg tgttccgtta attttgtaaa tactggctcg 2520
actgtaactc tcacatgtga caaagtatgg tttgttttgt tgggttttgt ttttgagaat 2580
ttttttgccc gtcccttttg tttcaaaagt ttcacgtctt ggtgcctttt gtgtgacacg 2640
55 ccttccgatg gcttgacatg cgcagatgtg agggacacgc tcaccttagc ctttaaggggg 2700
taggagtgat gtgttggggg aggccttgaga gcaaaaacga ggaagagggc tgagctgagc 2760
tttcggtctc cagaatgtaa gaagaaaaa ttttaacaaa aatctgaact ctcaaaagtc 2820
tatttttcta aactgaaaat gtaaatttat acatctattc aggagttgga gtgttggtgt 2880
tacctactga gtaggctgca gtttttgtat gttatgaaca tgaagttcat tattttgtgg 2940
60 ttttatttta ctttgtactt gtgtttgctt aaacaaagta acctgtttgg cttataaaca 3000

WO 00/42173

PCT/AU00/00011

5/8

cattgaatgc gctctattgc ccatgggata tgtgggtgtgt atccttcaga aaaattaaaa 3060
ggaaaaat 3068

5 <210> 12
<211> 4321
<212> DNA
<213> Rattus rattus

10 <400> 12
ccgcggagcc tcagctctac gcgcctggcg ccctccctac gcggggcgtec ccgaactcccg 60
cgcgcggttca ggctccgggt tgggaaccaa ggagggggag ggtgggtgcg ccgacccgga 120
aacaccatat aaggagcagg aaggatcccc cgccggaaca gacettatctt gggcagcgcc 180
ttatatggag tggcccaata tggccctgcc gcttccggct ctgggaggag gggcgaacgg 240
15 ggggttggggc gggggcaagc tgggaactcc aggagcctag cccgggaggg cactgccgct 300
gttccaatac taggctttcc aggagcctga gcgctcaggg tgccggagcc ggtcgcaggg 360
tggaagcgcc caccgctctt ggatgggagg tcttcacgct actccgggtc ctcccggtcg 420
gtccttccat attagggtct cctgcttccc atatatggcc atgtacgtca cggcggaggc 480
gggcccgtgc tgtttcagac ccttgaaata gaggccgatt cggggagtcg cgagagatcc 540
20 cagcgcgcag aacttgggga gccgcgcgcg cgattcgccg ccgcccgcag ctcccgccgc 600
cgcaagatcg gcccttgcgc cagcctccgc ggcagccctg cgtccaccac gggccgcggc 660
caccgccagc ctggggggccc acctacactc cccgcagtgt gcccctgcac cccgcagtga 720
acccggccaa catccggcga gtgtgcccctc agtagcttcc gcccggggct gcgcccacca 780
cccaacatca gctctccagc tcgcacgtcc gggatggcag cggccaaggc cgagatgcaa 840
25 ttgatgtctc cgtgcagat ctctgacccg ttccgctcct ttcctcactc acccaccatg 900
gacaactacc ccaaactgga ggagatgatg ctgctgagca acggggctcc ccagttcctc 960
ggtgctgccc gaaccccaga gggcagcggc ggcaataaca gcagcagcag cagcagcagc 1020
agcagcgggg gcggtggtgg gggcggcagc aacagcggca gcagcgcttt caatcctcaa 1080
ggggagccga gcgaacaacc ctacgagcac ctgaccacag gtaagcgggtg gtctgcgccg 1140
30 aggetgaatc ccccttcgtg actaccctaa cgtccagtc tttgcagcac ggacctgcac 1200
ctagatctta gggacgggat tgggatttcc ctctattcca cacagctcca gggacttggt 1260
ttagagggat gtctggggac cccccaaccc tccatccttg cgggtgcgcg gagggcagac 1320
cgtttgtttt ggatggagaa ctcaagttgc gtgggtgggt ggagtggggg agggtttggt 1380
ttgatgagca ggggtgcccc ctccccgcgc cgcgttgctg cgagccttgt ttgcagcttg 1440
35 tcccaagga agggctgaaa tctgtcacca gggatgtccc gccgcccagg gtaggggcgc 1500
gcattagctg tggccactag ggtgctggcg ggattccctc acccgggacg cctgctgcgc 1560
agcgtctcca gagctgcagt agagggggat tctctgtttg cgtcagctgt cgaaatggct 1620
ctgccactgg agcaggtcca ggaacattgc aatctgctgc tatcaattat taaccacatc 1680
gagagtcagt ggtagccggg cgacctcttg cctggccgct teggtctca tegtccagt 1740
40 attgctctcc agtaaccagg cctctctgtt ctctttcctg ccagagtcct tttctgacat 1800
cgctctgaat aacgagaagg cgctggtgga gacaagttat cccagccaaa ctaccgggtt 1860
gcctcccatc acctatactg gccgcttctc cctggagcct gcacccaaca gtggcaacac 1920
tttggtggcc gaaccccttt tcagcctagt cagtggcctt gtgagcatga ccaaccctcc 1980
aacctcttca tctcagcgc cttctccagc tgttctatcg tcttctctg cctcccagag 2040
45 cccacccctg agctgtgccc tgccgtccaa cgacagcagt cccatttact cagctgcacc 2100
cacctttcct actcccaaca ctgacatttt tcttgagccc caaagccagg cctttcctgg 2160
ctctgcaggc acagccttgc agtaccgcgc tctgacctac cctgccacca aggggtgggtt 2220
ccaggttccc atgatccctg actatctgtt tccacaacaa caggagagac tgagcctggg 2280
caccacagac cagaagccct tccagggtct ggagaaccgt accagcagc cttegcctac 2340
50 tccactatcc actatcaaag ccttcgccac tcagtcgggc tcccaggact taaaggctct 2400
taataacacc taccagtccc aactcatcaa accagccgc atgcgcaagt accccaaccg 2460
gccagcaag acacccccc atgaacgccc gtatgcttgc cctggttgagt cctgcgacgc 2520
ccgcttttct cgctcggatg agcttacacg ccacatccgc atccatacag gccagaagcc 2580
cttccagtgt cgaatctgca tgcgtaattt cagtctgagt gaccacctta ccaccacat 2640
55 ccgcacccac acaggcgaga agccttttgc ctgtgacatt tgtgggagaa agtttgccag 2700
gagtgatgaa cgcaagagge ataccaaaat ccacttaaga cagaaggaca agaaagcaga 2760
caaaagtgtc gtggcctcct cagctgcctc ttcctctctc tctaccat cccagtggc 2820
tacctcctac ccacccccgc ccaccacctc atttccatcc ccagtgcaca cctcttactc 2880
ctctccgggc tctctacct acccgtctcc tgcacacagt ggcttcccat cgcctcgggt 2940
60 ggccaccacc tatgcctccg tcccacctgc ttccctgccc caggtcagca ccttccagtc 3000

WO 00/42173

PCT/AU00/00011

6/8

tgcaggggtc agcaactcct tcagcacctc aacgggtcct tcagacatga cagcaacett 3060
 ttctcctagg acaattgaaa ttgctaaaag ggaatgaaag agagcaaagg gaggggagcg 3120
 cgagagacaa taaaggacag gaggggaagaa atggcccgcga agaggggctg cctcttaggt 3180
 cagatggaag atctcagagc caagtccttc tagtcagtag aaggcccgtt ggccaccagc 3240
 5 cctttcactt agcgtccctg cccctcccag tcccggctct tttgacttca gctgcctgaa 3300
 acagccacgt ccaagttctt cacctctatc caaaggactt gatttgcatt gtattggata 3360
 aaccatttca gcatcatctc caccacatgc ctggcccttg ctcccttcag cactagaaca 3420
 tcaagttggc tgaaaaaaa aatgggtctg ggccctcaga accctgccct gtatctttgt 3480
 acagcatctg tgccatggat tttgttttcc ttgggggtatt cttgatgtga agataatttg 3540
 10 catactctat tgtactatct ggagttaaata tctcactttg ggggaggggg agcaaagcca 3600
 agcaaaccac tggatgatcct ctattttgtg atgactcctg tgtgacatta ggtttgaaac 3660
 tttttttttt ttttgaagca gcagtcctag gtattaaact gagcatgtgt cagagtgttg 3720
 ttccgttaat tttgtaaata ctgctcgact gtaactctca catgtgacaa aatacggttt 3780
 gtttggttgg gttttttgtt gtttttgaaa aaaaaatttt ttttttgccc gtcccttttg 3840
 15 tttcaaaagt ttcacgtctt ggtgcctttg tgtgacacac cttgccgatg gctggacatg 3900
 tgcaatcgtg aggggacacg ctcacctcta gccttaaggg ggtaggagtg atgtttcagg 3960
 ggaggcttta gagcacgatg aggaagaggg ctgagctgag ctttggttct ccagaatgta 4020
 agaagaaaaa tttaaaacaa aaatctgaac tctcaaaagt ctattttttt aactgaaaat 4080
 gtagatttat ccatgttcgg gagttggaat gctgcggtta cctactgagt aggcggtgac 4140
 20 ttttgtatgc tatgaacatg aagttcatta ttttgtggtt ttattttact tcgtacttgt 4200
 gtttgcttaa acaaagtgc ttgtttggct tataaacaca ttgaatgcgc tttactgcc 4260
 atgggatatg tgggtgtgat ccttcagaaa aattaaaagg aaaataaaga aactaactgg 4320
 t 4321

 25
 <210> 13
 <211> 19
 <212> RNA
 <213> Rattus rattus
 30
 <400> 13
 acguccggga uggcagcgg 19

 35
 <210> 14
 <211> 19
 <212> RNA
 <213> Homo sapiens
 40
 <400> 14
 ucguccagga uggccgcgg 19

 45
 <210> 15
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 50 <221> misc_feature
 <222> (33)..(34)
 <223> 3'-3-linked T

 <220>
 55 <223> Description of Artificial Sequence: DNAzyme

 <400> 15
 caggggacag gctagctaca acgacgttgc gggt 34
 60

WO 00/42173

PCT/AU00/00011

7/8

5 <210> 16
 <211> 34
 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc_feature
 <222> (33)..(34)
 <223> 3'-3-linked T
 10 <220>
 <223> Description of Artificial Sequence: DNAzyme
 <400> 16
 15 tgcaggggag gctagctaca acgaaccggt gcgt 34
 <210> 17
 <211> 34
 20 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> misc_feature
 25 <222> (33)..(34)
 <223> 3'-3-linked T
 <220>
 <223> Description of Artificial Sequence: DNAzyme
 30 <400> 17
 catcctggag gctagctaca acgagagcag gctt 34
 35 <210> 18
 <211> 34
 <212> DNA
 <213> Artificial Sequence
 40 <220>
 <221> misc_feature
 <222> (33)..(34)
 <223> 3'-3-linked T
 45 <220>
 <223> Description of Artificial Sequence: DNAzyme
 <400> 18
 50 tcagctgcag gctagctaca acgactcggc cttt 34
 <210> 19
 <211> 34
 <212> DNA
 55 <213> Artificial Sequence
 <220>
 <221> misc_feature
 <222> (33)..(34)
 60 <223> 3'-3-linked T

WO 00/42173

PCT/AU00/00011

8/8

<220>

<223> Description of Artificial Sequence: DNAzyme


5 <400> 19
gcggggacag gctagctaca acgacagctg catt 34

10 <210> 20
<211> 15
<212> DNA
<213> Rattus rattus

15 <400> 20
cttggccgct gccat 15

20

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☒ Page(s) _____ of Express Mail were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ *Scanned copy is best available.*